

**A peer-reviewed version of this preprint was published in PeerJ on 6 July 2018.**

[View the peer-reviewed version](https://doi.org/10.7717/peerj.5212) (peerj.com/articles/5212), which is the preferred citable publication unless you specifically need to cite this preprint.

Taylor MP. 2018. *Xenoposeidon* is the earliest known rebbachisaurid sauropod dinosaur. PeerJ 6:e5212 <https://doi.org/10.7717/peerj.5212>

# ***Xenoposeidon* is the earliest known rebbachisaurid sauropod dinosaur**

**Michael Taylor** <sup>Corresp. 1</sup>

<sup>1</sup> Department of Earth Sciences, University of Bristol

Corresponding Author: Michael Taylor  
Email address: dino@miketaylor.org.uk

*Xenoposeidon proneneukos* is a sauropod dinosaur represented by a single partial dorsal vertebra, NHMUK R2095, which consists of the centrum and the base of a tall neural arch. Despite its fragmentary nature, it is recognisably distinct from all other sauropods, and is here diagnosed with five unique characters. One character previously considered unique is here recognised as shared with *Rebbachisaurus garasbae*: an “M”-shaped arrangement of laminae on the lateral face of the neural arch. Following the more complete *Rebbachisaurus garasbae*, these laminae are now interpreted as ACPL and lateral CPRL, which intersect anteriorly; and PCDL and CPOL, which intersect posteriorly. Similar arrangements are also seen in some other rebbachisaurid specimens (though not all, possibly due to serial variation), but never in non-rebbachisaurid sauropods. *Xenoposeidon* is therefore referred to Rebbachisauridae. Due to its elevated parapophysis, the holotype vertebra is considered a posterior dorsal despite its elongate centrum. Since *Xenoposeidon* is from the from the Berriasian–Valanginian (earliest Cretaceous) Ashdown Beds Formation of the Wealden Supergroup of southern England, it is the earliest known rebbachisaurid by some 10 million years. Electronic 3D models were invaluable in determining *Xenoposeidon*'s true affinities: descriptions of complex bones such as sauropod vertebrae should always provide them where possible.

# 1 *Xenoposeidon* is the earliest known rebbachisaurid 2 sauropod dinosaur

3 Michael P. Taylor. Department of Earth Sciences, University of Bristol, Bristol BS8 1RJ,  
4 England.  
5 [dino@miketaylor.org.uk](mailto:dino@miketaylor.org.uk)

## 6 Abstract

7 *Xenoposeidon proneneukos* is a sauropod dinosaur represented by a single partial dorsal vertebra,  
8 NHMUK R2095, which consists of the centrum and the base of a tall neural arch. Despite its  
9 fragmentary nature, it is recognisably distinct from all other sauropods, and is here diagnosed  
10 with five unique characters. One character previously considered unique is here recognised as  
11 shared with *Rebbachisaurus garasbae*: an “M”-shaped arrangement of laminae on the lateral face  
12 of the neural arch. Following the more complete *Rebbachisaurus garasbae*, these laminae are  
13 now interpreted as ACPL and lateral CPRL, which intersect anteriorly; and PCDL and CPOL,  
14 which intersect posteriorly. Similar arrangements are also seen in some other rebbachisaurid  
15 specimens (though not all, possibly due to serial variation), but never in non-rebbachisaurid  
16 sauropods. *Xenoposeidon* is therefore referred to Rebbachisauridae. Due to its elevated  
17 parapophysis, the holotype vertebra is considered a posterior dorsal despite its elongate centrum.  
18 Since *Xenoposeidon* is from the from the Berriasian–Valanginian (earliest Cretaceous) Ashdown  
19 Beds Formation of the Wealden Supergroup of southern England, it is the earliest known  
20 rebbachisaurid by some 10 million years. Electronic 3D models were invaluable in determining  
21 *Xenoposeidon*'s true affinities: descriptions of complex bones such as sauropod vertebrae should  
22 always provide them where possible.

## Table of Contents

Introduction.....	2
Anatomical Abbreviations.....	2
Institutional Abbreviations.....	3
Reinterpretation.....	3
Serial position.....	6
Revised Reconstruction.....	7
Systematic Palaeontology.....	7
Discussion.....	8
Age.....	8
Wealden Rebbachisaurids.....	9
3D models of complex bones.....	9
Acknowledgements.....	9
References.....	10
Figure Captions.....	12
Supplementary Files.....	13

## 23 Introduction

24 *Xenoposeidon proneneukos* is a neosauropod sauropod dinosaur from the Berriasian–Valanginian  
25 (earliest Cretaceous) Ashdown Beds Formation of the Wealden Supergroup of southern England.  
26 It is represented by a single partial mid-to-posterior dorsal vertebra, NHMUK R2095 (BMNH  
27 R2095 at the time of the original description by Taylor and Naish 2007). This element consists of  
28 the centrum and the base of a tall neural arch, broken off below the transverse processes and  
29 zygapophyses. Despite its fragmentary nature, it is recognisably different from all other  
30 sauropods, and Taylor and Naish (2007) diagnosed it on the basis of six characters that they  
31 considered unique among sauropods.

32 D’Emic (2012:651) asserted that “the absence of diagnostic features renders *Xenoposeidon* a  
33 nomen dubium”. However, his assessment was mistaken in several respects. For example, the  
34 extension of the base of the neural arch to the posterior extremity of the centrum is clearly not, as  
35 he asserted, due to damage. D’Emic claimed that dorsal vertebrae illustrated by Osborn and  
36 Mook (1921:plates LXIX and LXXII) have forward-sloping neural arches resembling those of  
37 *Xenoposeidon*: in reality, only one posterior dorsal vertebrae out of four complete dorsal columns  
38 illustrated in that monograph shows a forward slope, and it differs so much from its fellows that  
39 this can only be interpreted as the result of crushing. D’Emic further claimed that the lamina  
40 patterns observed in *Xenoposeidon* can be recognised in other sauropods, but I have been unable  
41 find morphology resembling them in the descriptions he suggests: Osborn and Mook 1921 for  
42 *Camarasaurus*, Riggs 1903 for *Brachiosaurus* (probably a typo for Riggs 1904, which also does  
43 not depict similar patterns), Carballido et al. 2011 for *Tehuelchesaurus*. A similar pattern does  
44 appear in *Rebbachisaurus*, as will be discussed below. D’Emic (2012:651) is probably correct  
45 that the “asymmetric neural canal” described by Taylor and Naish (2007:1553–1554) is a  
46 misreading of the tall centroprezygapophyseal fossae as being the anterior portion of the neural  
47 canal: as Taylor and Naish pointed out, “The vacuity is filled with matrix, so the extent of its  
48 penetration posteriorly into the neural arch cannot be assessed”. Nevertheless, the shape and size  
49 of the fossa is unique among sauropods, and it is bounded by laminae which do not seem to be  
50 medial CPRLs. In summary, *Xenoposeidon proneneukos* is a valid, diagnosable taxon, *contra*  
51 D’Emic (2012).

52 Taylor and Naish (2007:1554–1557) compared the *Xenoposeidon* vertebra to those of the main  
53 neosauropod groups — Diplodocoidea, Camarasauridae, Brachiosauridae and Titanosauria —  
54 and concluded that it could not be convincingly referred to any of these groups. Their  
55 phylogenetic analysis (pp. 1157–1558 and figure 6) corroborated this by recovering  
56 *Xenoposeidon* as a neosauropod in all most parsimonious trees, but in a polytomy with all other  
57 neosauropods, wholly unresolved save that the clade Flagellicaudata was preserved in all MPTs.

58 In light of Wilson and Allain’s (2015) redescription of *Rebbachisaurus garasbae*, and the  
59 availability of more photographs and models of rebbachisaurid material, it has now become  
60 possible to reinterpret the idiosyncratic system of laminae found in *Xenoposeidon*, and to refer it  
61 confidently to an existing family-level clade.

## 62 Anatomical Abbreviations

- 63 • aEI — average elongation index *sensu* Chure et al. 2010: length of a centrum divided by  
64 the average of the height and width of the posterior articular surface.

- 65 • ACPL — anterior centroparapophyseal lamina.
- 66 • CPOL — centropostzygapophyseal lamina.
- 67 • CPRF — centroprezygapophyseal fossa.
- 68 • CPRL — centroprezygapophyseal lamina.
- 69 • EI — elongation index *sensu* Wedel et al. 2000: length of a centrum divided by the height
- 70 of the posterior articular surface.
- 71 • PCDL — posterior centrodiapophyseal lamina.
- 72 • PCPL — posterior centroparapophyseal lamina.
- 73 • POSL — postspinal lamina.
- 74 • Postzyg — postzygapophysis.
- 75 • PPDL — paradiapophyseal lamina.
- 76 • Prezyg — prezygapophysis.
- 77 • PRPL — prezygaparapophyseal lamina.
- 78 • PRSL — prespinal lamina.
- 79 • SDL — spinodiapophyseal lamina.

## 80 Institutional Abbreviations

- 81 • IWCMS — Isle of Wight County Museum Service at Dinosaur Isle, Sandown, Isle of
- 82 Wight, England.
- 83 • MIWG — Museum of Isle of Wight Geology (now Dinosaur Isle Visitor Centre),
- 84 Sandown, Isle of Wight, England.
- 85 • MNHN — Muséum National d'Histoire Naturelle, Paris, France.
- 86 • NHMUK — the Natural History Museum, London, England.
- 87 • NMC — Canadian Museum of Nature (previously National Museum of Canada), Ottawa,
- 88 Ontario, Canada.
- 89 • “WN” — “without number”, an informal designation for specimens awaiting accession.

## 90 Reinterpretation

91 Taylor and Naish’s (2007) history, geography, geology and description of the *Xenoposeidon*  
92 specimen requires no revision, and should continue to be considered definitive: this paper does  
93 not supersede it, but should be read in conjunction with it.

94 The illustrations of the specimen in the original paper, however, were in monochrome and  
95 omitted the dorsal and ventral views. The present paper supplements these illustrations with a  
96 colour depiction from all six cardinal directions (Figure 1), and a high-resolution 3D model of the  
97 specimen (supplementary file AA).

98 More importantly, Taylor and Naish’s (2007) interpretation of some features of the vertebra,  
99 particularly the “M”-shaped complex of laminae on the lateral faces of the neural arch, was  
100 mistaken. Although the neural spine and dorsal part of the neural arch are missing, including the  
101 pre- and postzygapophyses and lateral processes, they wrote that “sufficient laminae remain to  
102 allow the positions of the processes to be inferred with some certainty”. But their inferences were  
103 incorrect. Taylor and Naish (2007:1553) interpreted the cross-shaped structure on the  
104 anterodorsal part of the left lateral face of the neural arch as the site of the parapophysis, despite  
105 the lack of any articular facet in that location. This influenced their interpretation of the four  
106 laminae that met at that point as the ACPL below, the PPDL above, the PRPL anteriorly and an

107 unnamed accessory infraparapophyseal lamina posteroventrally, which they interpreted as  
108 homologous with a PCPL (Figure 2A). Similarly, they did not attempt to identify either the long  
109 lamina running up the posterior edge of the lateral face of the neural arch (designating it only  
110 “posterior lamina”) or the lamina forming a shallow “V” with the “accessory infraparapophyseal  
111 lamina”, simply calling it an “accessory postzygapophyseal lamina” (Figure 2A)

112 Among the various unusual features of the *Xenoposeidon* vertebra, the “M”-shaped set of laminae  
113 is immediately apparent in lateral view (Figure 3A): a line can be traced from the anterior margin  
114 of the neural arch’s lateral face up the ACPL to the cross that was interpreted as the parapophysis,  
115 then posteroventrally down the “accessory infraparapophyseal lamina”, then posterodorsally up  
116 the “accessory postzygapophyseal lamina” and finally down the posterior margin of the neural  
117 arch’s lateral face, along the “posterior lamina”. Photographs of other specimens that were  
118 available to us at this time did not apparently manifest similar features.

119 But subsequent work on *Rebbachisaurus garasbae* (Wilson 2012:100, Wilson and Allain 2015)  
120 — and an associated video of the rotating vertebra (see acknowledgements) — show that  
121 *Rebbachisaurus* has a similar complex of laminae (Figure 3B), which are described by Wilson  
122 and Allain (2015:6) as the second of the eight autapomorphies that they listed for the species:  
123 “infrazygapophyseal laminae (lat. CPRL, CPOL) that intersect and pass through neighbouring  
124 costal laminae (ACPL, PCDL) to form an ‘M’ shape”.

125 Because the illustrated dorsal vertebra of *Rebbachisaurus* — MNHN MRS 1958 — is  
126 substantially complete, it is possible to follow the trajectories of the laminae that participate in  
127 the “M” to their apophyses, and so determine their true identities. The two vertically oriented  
128 laminae — the outer pillars of the “M” — continue up past the top of the “M”. The anterior one  
129 supports the parapophysis, and the posterior supports the diapophysis. And the two laminae that  
130 form the valley in the middle of the “M” support the prezygapophyses and postzygapophyses: in  
131 both cases, as noted by Wilson and Allain, they intersect the vertical lamina before continuing to  
132 meet their respective zygapophyses. The four laminae that make up the “M”, from anterior to  
133 posterior, are therefore the ACPL, posterior part of the lateral CPRL, anterior part of the CPOL  
134 and PCDL. Of these, the intersection between the ACPL and lateral CPRL is clearly visible in left  
135 lateral view of MNHN MRS 1958. The intersection between the CPOL and PCDL is less  
136 apparent in this view, though clear in three dimensions. Both laminae continue dorsally beyond  
137 this intersection, but their paths are somewhat changed at the point of contact, with the dorsal  
138 portion of the PCDL inclining more anteriorly, and the rod-like CPOL apparently passing through  
139 the sheet of bone formed by the PCDL to meet the postzygapophysis.

140 The referred *Rebbachisaurus garasbae* specimen NMC 50844 described and illustrated by  
141 Russell (1996:388–390 and figure 30) is also broadly consistent with this morphology. It is not  
142 possible to be definite about the laminar intersection based only on line drawings of the specimen  
143 from the four cardinal directions, but, as illustrated in Russell’s figure 30c, the lateral CPRL does  
144 appear to pass through the ACPL. The CPOL seems in this specimen to originate posterior to the  
145 PCDL, not intersecting with it. But this difference from the holotype dorsal may be serial  
146 variation since, as Russell notes, the relatively longer centrum of his specimen indicates a more  
147 anterior serial position than for the holotype’s dorsal vertebra; and this interpretation is  
148 corroborated by the observation that, based on lamina trajectories, the anteroposterior distance  
149 between the parapophysis and diapophysis was less in NMC 50844 than in the holotype.

150 In light of these *Rebbachisaurus* specimens, the mysterious laminae of *Xenoposeidon* are easily  
151 explained. It is now apparent that the cross on the side of the *Xenoposeidon* vertebra is not the  
152 site of the parapophysis, as Taylor and Naish (2007:1553) proposed, but merely the intersection

153 of two laminae that pass right through each other: the ACPL, running dorsolaterally, and the  
154 lateral CPRL, extending anterodorsally to the (missing) prezygapophysis (Figure 2B). Similarly,  
155 the “posterior lamina” is the PCDL, and it intersects with the CPOL, though the intersection is  
156 lost in NHMUK R2095 (Figure 2B). Both the parapophysis and diapophysis of the *Xenoposeidon*  
157 vertebrae would have been located some distance above the preserved portion, the former  
158 anterior to the latter.

159 It appears from Dalla Vecchia (1999:figure 47, left part) that in the holotype and only vertebra of  
160 *Histriasaurus boscarollii*, “WN-V6”, the CPOL on the right side of the vertebra intersects with  
161 the PCDL in the same way as in *Rebbachisaurus*, though it is not possible to determine whether  
162 the lateral CPRL similarly intersects the ACPL. Dorsal vertebrae of other rebbachisaurid  
163 sauropods, however, do not appear to feature the distinctive “M” and intersecting laminae of  
164 *Rebbachisaurus* and *Xenoposeidon*:

- 165 • The 3D model of a dorsal vertebra of *Nigersaurus* (Sereno et al. 2007) shows that the  
166 lateral CPRLs originate anterior to the ACPLs and the CPOLs posterior to the PCDLs, so  
167 that there is no intersection. A subtle “V” shape does appear high up on the lateral faces of  
168 the neural arch, between the ACPL and the PCDL, but it seems unrelated to the lateral  
169 CPRL and CPOL.
- 170 • Unpublished 3D models of an anterior dorsal neural arch and a more posterior dorsal  
171 vertebra of *Katpensaurus* (pers. comm., Lucio M. Ibiricu) as illustrated in figures 3A and  
172 5A of Ibiricu at el. (2017) show that in both vertebrae, the lateral CPRLs originate anterior  
173 to the ACPLs, and the CPOLs seem to originate posterior to the PCDLs — though  
174 damage to the posterior portion makes the latter uncertain.
- 175 • The laminae do not appear to intersect in the illustrated dorsal vertebra of  
176 *Demandasaurus* (Fernández-Baldor et al. 2011:figure 9).
- 177 • The sole known vertebra of *Nopcsaspondylus* seems to have an entirely different pattern  
178 of lamination (Mannion 2010:figure 5) with no lamina intersections like those of MNHN  
179 MRS 1958.

180 No determination can be made for other rebbachisaurids as they are insufficiently preserved (e.g.  
181 *Limaysaurus*, *Amazonsaurus*), or illustrated (e.g. *Cathartesaura*), or simply lack posterior dorsal  
182 vertebral material (e.g. *Rayososaurus*, *Tataouinea*, *Comahuesaurus*, *Zapalasaurus*).

183 However, we cannot rule out the possibility that complete and well-preserved posterior dorsal  
184 vertebrae of most or all rebbachisaurids have *Rebbachisaurus*-like intersecting laminae: even in  
185 those species for which a well-preserved vertebra lacks them, this could be due to serial variation,  
186 with these features only fully developing in the most posterior dorsals.

187 *Xenoposeidon*, then, resembles *Rebbachisaurus* in the possession of a distinctive “M” on the  
188 lateral face of the neural arch, in the intersecting lateral CPRL and ACPL, and in the elevation of  
189 the parapophysis above the level of the prezygapophysis — a complex of related features.  
190 Although at first glance they appear rather different, *Xenoposeidon* and *Rebbachisaurus*, while  
191 geometrically different, are topologically similar.

192 Regarding the significance of the elevated parapophysis, since no complete or nearly complete  
193 rebbachisaurid dorsal column has been described, comparisons with other, better represented  
194 sauropods are warranted. In the probable basal diplodocoid *Haplocanthosaurus*, the dorsal  
195 margin of the parapophyseal facet reaches the level of, and is coincident with, the  
196 prezygapophyseal facet around dorsal vertebra 7 or 8, but never rises any higher than this in more

197 posterior vertebrae (Hatcher 1903:plate I). In the more distantly related diplodocid diplodocoids  
198 *Apatosaurus* and *Diplodocus*, the parapophysis never migrates far enough dorsally to reach a  
199 position level with the prezygapophyses, even in the most posterior dorsals (Gilmore 1936:plate  
200 XXV; Hatcher 1901:plates VII, VIII).

201 Taylor and Naish (2007:1554) argued that *Xenoposeidon* could not at that time be convincingly  
202 referred to Rebbachisauridae because *Rebbachisaurus* differs from NHMUK R2095 in five ways:  
203 “possession of a very prominent PCPL, large and laterally diverging prezygapophyses,  
204 depressions at the base of the neural arch (Bonaparte 1999:173), lateral foramina not set within  
205 fossae, and a strongly arched ventral border to the centrum.” Of these features, the first is now  
206 recognised as occurring in *Xenoposeidon*; the second appears to be an outright error, as the  
207 prezygapophyses of *Rebbachisaurus* meet on the midline, and in any case the situation in  
208 *Xenoposeidon* is not known. “Depressions at the base of the neural arch” seems to be a  
209 mistranslation of Bonaparte’s original Spanish, “profundas depresiones en la base de la espina  
210 neural”, which refers not to the neural arch but the neural spine, and since this portion is not  
211 preserved in *Xenoposeidon*, it is not informative for our purposes. The 3D model of the  
212 *Rebbachisaurus* dorsal shows that in fact its lateral foramina are set in shallow depression,  
213 similar in quality if not in degree to those of *Xenoposeidon*. This leaves the stronger arching of  
214 the ventral border of the centrum in *Rebbachisaurus*, a feature that in isolation is not convincing.

215 In conclusion, the weight of morphological evidence supports including *Xenoposeidon* within  
216 Rebbachisauridae. This is in accordance with the observation of Taylor and Naish (2007:1557), in  
217 whose phylogenetic analysis “various most-parsimonious trees also recover *Xenoposeidon* in  
218 many other positions, including as a ... rebbachisaurid.”

## 219 Serial position

220 The serial position of the *Rebbachisaurus garasbae* holotype dorsal vertebra MNHN MRS 1958  
221 is not definitely known. However, it has been uniformly referred to as a posterior dorsal, most  
222 likely due to the very elevated position of its parapophyses and Lavocat’s (1954) initial  
223 assessment of it as “une des dernières dorsales” (one of the last dorsals) — perhaps made with  
224 knowledge of the spatial relation of bones in the quarry.

225 The position of the *Xenoposeidon proneneukos* holotype vertebra NHMUK R2095 is of course  
226 even more difficult to determine in light of the limited nature of the specimen, though its  
227 similarity to MNHN MRS 1958 suggests a similar position. Taylor and Naish (2007:1553) wrote  
228 that “the high position of the parapophysis on the neural arch of R2095 indicates a mid to  
229 posterior placement of the vertebra within the dorsal column, but, because the prezygapophyses  
230 must have been dorsal to it, it was probably not among the most posterior vertebrae in the  
231 sequence.” With the location of the parapophysis now interpreted as significantly higher than  
232 previously thought, and probably well above the prezygapophysis, an even more posterior  
233 position is indicated.

234 This posterior serial position is surprising in light of the anteroposterior length of the  
235 *Xenoposeidon* centrum. Its posterior articular surface measures 160 mm high by 170 mm wide,  
236 while the length of even the preserved portion of the centrum is 190 mm, and it must have been at  
237 least 200 mm long when complete (Taylor and Naish 2007:table 1). As noted by Taylor and Naish  
238 (2007:1554), “the length of the centrum, especially in so posterior a dorsal vertebra, argues  
239 against [a diplodocoid identity]: the posterior dorsal centra of diplodocoids typically have  $EI <$   
240 1.0, compared with 1.25 for R2095” — or 1.21 using the aEI of Chure et al. (2010:384).  
241 However, rebbachisaurids may be unusual among diplodocoids in this respect — perhaps



242 unsurprisingly, as they diverged early from the line leading to diplodocids, with their  
243 characteristically short dorsal centra, and likely retained something more similar to the ancestral  
244 neosauropod condition. Wilson and Allain (2015:8) give the centrum measurements of MNHN  
245 MRS 1958 as posterior height 231 mm, posterior width 220 mm and length 220 mm. This yields  
246 an aEI of 0.98, meaning that the *Xenoposeidon* centrum is only 24% more elongate than that of  
247 *Rebbachisaurus*. This is a significant difference, but not an outlandish one. For comparison, the  
248 centrum of the basal rebbachisaurid *Histriasaurus boscarollii* holotype “WN-V6” is relatively  
249 elongate, with its posterior articular surface measuring 150 mm high and centrum length of “more  
250 than 200 mm” (Dalla Vecchia 1998:122) yielding an EI of > 1.33. Also, the aEIs of the last four  
251 dorsal vertebrae of the *Brachiosaurus altithorax* holotype FMNH PR 25107 are 1.34, 1.27, 1.19  
252 and 0.96 (calculated from the table of Riggs 1904:34): so aEIs of sauropod dorsals can vary,  
253 within two serial positions of the same individual, from values below that of MNHN MRS 1958  
254 to above that of NHMUK R2095.

255 In conclusion, while the evidence regarding the serial position of NHMUK R2095 remains  
256 equivocal, it suggests a more posterior position than previous inferred — it can be fairly  
257 confidently described as “posterior” rather than “mid-to-posterior” — but it is unlikely to be the  
258 very last dorsal.

## 259 Revised Reconstruction

260 In light of the reassignment of *Xenoposeidon* to Rebbachisauridae, and the reinterpretation of its  
261 laminae, I present a new reconstruction of how the vertebra NHMUK R2095 might have looked  
262 when complete (Figure 4). As in MNHN MRS 1958, the parapophysis and diapophysis are both  
263 elevated above the zygapophyses. The lateral CPRL and ACPL meet at a point where they  
264 project outwards about the same distance from the vertebra, as is apparent from the preserved  
265 portion of the vertebra; but the CPOL is assumed to pass through a sheet-like PCDL as in  
266 *Rebbachisaurus*, because it is clear from breakage in NHMUK R2095 that the PCDL extended  
267 further from the body of the neural arch than the preserved portion indicates. The neural spine,  
268 composed as in *Rebbachisaurus* of pre- and post-spinal laminae together with the left and right  
269 SDLs, is shown fading out at the top, as there is no way to determine its height. The condyle that  
270 is the centrum’s anterior articular surface is reconstructed as only slightly convex, as in  
271 *Rebbachisaurus*.

272 It is instructive to compare this with the original reconstruction of the vertebrae (Taylor and  
273 Naish:figure 5). The new reconstruction has a taller neural arch, a far more elevated  
274 parapophysis, a more posteriorly located diapophysis (no longer dorsal to the parapophysis) and a  
275 shallower condyle, as that of the original reconstruction was drawn with those of brachiosaurs in  
276 mind.

## 277 Systematic Palaeontology

278 Dinosauria Owen, 1842  
279 Saurischia Seeley, 1888  
280 Sauropodomorpha Huene, 1932  
281 Sauropoda Marsh, 1878  
282 Neosauropoda Bonaparte, 1986  
283 Rebbachisauridae Sereno et al., 1999  
284 *Xenoposeidon* Taylor and Naish, 2007

285 *Xenoposeidon proneneukos* Taylor and Naish, 2007

286 **Holotype.** NHMUK R2095, the Natural History Museum, London. A mid posterior dorsal  
287 vertebra consisting of partial centrum and neural arch.

288 **Revised diagnosis:** Differs from all other sauropods in the following characters:

- 289 1. neural arch covers dorsal surface of centrum, with its posterior margin continuous with  
290 that of the centrum;
- 291 2. neural arch slopes anteriorly 35 degrees relative to the vertical;
- 292 3. broad, flat area of featureless bone on lateral face of neural arch;
- 293 4. very large, teardrop-shaped centroprezygapophyseal fossa.
- 294 5. arched laminae form vaulted boundary of centroprezygapophyseal fossa.

295 The “arched laminae” of #5 are not the medial CPRLs, as these arise from the neural arch  
296 pedicels — and the laminae arising from the pedicels cannot instead be regarded lateral CPRLs,  
297 as those laminae are located on the lateral face of the neural arch, intersecting with the ACPLs.  
298 Furthermore, the point where the supporting laminae meet at the top of their arch is located some  
299 way posterior to the inferred location of the prezygapophyses (Figure 5).

## 300 Discussion

### 301 Age

302 As shown by the Wilson and Allain (2015:table 1), the 19 then-recognised rebbachisaurids (of  
303 which 13 had been named) span the middle third of the Cretaceous. The earliest recognised taxon  
304 is *Histriasaurus boscarollii* from the upper Hauterivian or lower Barremian limestones of  
305 southwest Istria, Croatia. Seven taxa, of which five are named, survived at least to the  
306 Cenomanian (earliest Late Cretaceous), of which two (*Katepensaurus goicoecheai* and  
307 *Limaysaurus tessonei*) may be from the Turonian age.

308 As discussed by Taylor and Naish (2007:1547–1548), the precise location and horizon where  
309 NHMUK R2095 was excavated was not recorded in the specimen’s original brief description,  
310 which only said “the Wealden of Hastings” (Lydekker 1893:276). However, records of the  
311 collection of Philip James Rufford, who collected the specimen, indicate that the most likely  
312 location is Ecclesbourne Glen, a mile or two east of Hastings, East Sussex (see discussion in  
313 Taylor and Naish 2007:1548). The units exposed at Ecclesbourne Glen are part of the Ashdown  
314 Beds Formation, which straddles the Berriasian/Valanginian boundary; but the part of the  
315 formation at that location is from the earlier Berriasian age. If this assessment is correct, then  
316 *Xenoposeidon* is from the very earliest Cretaceous, giving it an age of around 140 million years  
317 — about 10 million years earlier than *Histriasaurus*.

318 This early age is consonant with a basal position within Rebbachisauridae, a possibility that is  
319 corroborated by *Xenoposeidon*’s camerate internal morphology compared with the camellate  
320 centra of most rebbachisaurids. However, further material will be required before numerical  
321 phylogenetic work can firmly establish its position within the group.

## 322 Wealden Rebbachisaurids

323 Although *Xenoposeidon* is the first named Rebbachisaurid from the Wealden Supergroup of  
324 southern England, other material from this unit has been referred to Rebbachisauridae. Naish and  
325 Martill (2001:plate 36, opposite page 236) illustrated some isolated sauropod teeth  
326 IWCMS.2001.201–203, and these were referred to Rebbachisauridae by Sereno and Wilson  
327 (2005:174). Mannion (2009) described a partial rebbachisaurid scapula MIWG 6544. Finally,  
328 Mannion et al. (2011) described a proximal caudal neural arch MIWG 5384, which they also  
329 interpreted as rebbachisaurid. All of these specimens are from the Barremian Wessex Formation  
330 of the Isle of Wight, so they could all belong to the same species or genus. However, since the  
331 likely Berriasian age of NHMUK R2095 makes it 10–15 Mya older than these specimens, it is  
332 unlikely that they belong to *Xenoposeidon*, but to some other as yet-unnamed rebbachisaurid.  
333 Thus it is likely that the Wealden Supergroup contains at least two rebbachisaurid sauropods.

## 334 3D models of complex bones

335 Electronic 3D models were invaluable in determining *Xenoposeidon*'s true affinities. Most  
336 obviously, the model of the *Xenoposeidon* vertebra itself, created by Heinrich Mallison, has  
337 functioned as an invaluable proxy for the fossil itself when I am unable to visit the NHMUK, and  
338 I have consulted it many times in writing this paper. I would also have been unable to determine  
339 to my own satisfaction whether the *Katepensaurus* dorsals feature intersecting laminae like those  
340 of *Rebbachisaurus* without the models provided by Lucio M. Ibiricu. Although no true model is  
341 available for the *Rebbachisaurus* dorsal itself or for the dorsal vertebrae of *Nigersaurus*, rotating  
342 videos were crucial in enabling me to understand their morphology. When interpreting specimens  
343 for which no such models exist, such as Russell's (1996) referred *Rebbachisaurus* specimen  
344 NMC 50844, the conclusions reached using only 2D representations — whether photographs or  
345 drawings — are much less well founded.

346 Techniques such as photogrammetry (see e.g. Falkingham 2012; Mallison and Wings 2014) are  
347 reducing the barriers to the creation of high-quality 3D models in full colour. Doing so is now  
348 inexpensive in both time and money. In light of our discipline's goal of making palaeontology  
349 more accessible and reproducible, then, it should become increasingly routine in the 21st Century  
350 to provide 3D models as a standard part of the description of complex bones such as sauropod  
351 vertebrae.

## 352 Acknowledgements

353 I thank Sandra D. Chapman (Natural History Museum, London) for access to the *Xenoposeidon*  
354 specimen, and Heinrich Mallison (Palaeo3D) who went far beyond the call of duty in building the  
355 3D model of NHMUK R2095 and talking me through aspects of photogrammetry. I am also  
356 grateful to Jeff Wilson (University of Michigan) and Ronan Allain (Muséum National d'Histoire  
357 Naturelle, Paris) for sharing high-resolution photographs of the French *Rebbachisaurus* vertebra,  
358 and to Mathew J. Wedel (Western University of Health Sciences) and Darren Naish (University  
359 of Southampton) for helpful discussion. Lucio M. Ibiricu kindly provided access to unpublished  
360 3D models of an anterior dorsal neural arch and a more posterior dorsal vertebra of  
361 *Katepensaurus*.

362 As noted in Taylor (2015), this project began when I recognised the true identity of the curious  
363 laminae on the *Xenoposeidon* vertebra while viewing a rotating video of the *Rebbachisaurus*  
364 *garasbae* holotype dorsal vertebra MNHN MRS 1958 on the University of Michigan Museum of

365 Paleontology's UMORF web-site (University of Michigan Online Repository of Fossils) at  
366 <https://umorf.ummp.lsa.umich.edu/wp/gallery/vertebrate-animations/>. This video was based on a  
367 3D reconstruction created from CT scans performed at the AST-RX (Accès Scientifique à la  
368 Tomographie à Rayons X) of the MNHN by F. Goussard.

## 369 References

- 370 Bonaparte, Jose F. 1986. Les dinosaures (Carnosaures, Allosauridés, Sauropodes, Cétiosauridés)  
371 du Jurassique moyen de Cerro Cónдор (Chubut, Argentina). *Annales de Paléontologie* **72**:325–  
372 386.
- 373 Bonaparte, Jose F. 1999. Evolucion de las vertebras presacras en Sauropodomorpha [Evolution of  
374 the presacral vertebrae in Sauropodomorpha]. *Ameghiniana* **36**(2):115–87.
- 375 Carballido, Jose L., Oliver W. M. Rauhut, Diego Pol and Leonardo Salgado. 2011. Osteology and  
376 phylogenetic relationships of *Tehuelchesaurus benitezii* (Dinosauria, Sauropoda) from the  
377 Upper Jurassic of Patagonia. *Zoological Journal of the Linnean Society* **163**:605–662.  
378 doi:10.1111/j.1096-3642.2011.00723.x
- 379 Chure, Daniel, Brooks B. Britt, John A. Whitlock and Jeffrey A. Wilson. 2010. First complete  
380 sauropod dinosaur skull from the Cretaceous of the Americas and the evolution of sauropod  
381 dentition. *Naturwissenschaften* **97**(4):379–91. doi:10.1007/s00114-010-0650-6
- 382 Dalla Vecchia, Fabio M. 1998. Remains of Sauropoda (Reptilia, Saurischia) in the Lower  
383 Cretaceous (Upper Hauterivian/Lower Barremian) Limestones of SW Istria (Croatia).  
384 *Geologia Croatica* **51**(2):105–134.
- 385 Dalla Vecchia, Fabio M. 1999. Atlas of the sauropod bones from the Upper Hauterivian-Lower  
386 Barremian of Bale/Valle (SW Istria, Croatia). *Natura Nascosta* **18**:6–41.
- 387 D'Emic, Michael D. 2012. The early evolution of titanosauriform sauropod dinosaurs. *Zoological*  
388 *Journal of the Linnean Society* **166**:624–671.
- 389 Falkingham, Peter L. 2012. Acquisition of high resolution 3D models using free, open-source,  
390 photogrammetric software. *Palaeontologia Electronica* **15**(1):1T. 15 pages. [http://palaeo-](http://palaeo-electronica.org/content/issue1-2012technical-articles/92-3d-photogrammetry)  
391 [electronica.org/content/issue1-2012technical-articles/92-3d-photogrammetry](http://palaeo-electronica.org/content/issue1-2012technical-articles/92-3d-photogrammetry)
- 392 Fernández-Baldor, Fidel Torcida, José Ignacio Canudo, Pedro Huerta, Diego Montero, Xabier  
393 Pereda Suberbiola and Leonardo Salgado. 2010. *Demandasaurus darwini*, a new  
394 rebbachisaurid sauropod from the Early Cretaceous of the Iberian Peninsula. *Acta*  
395 *Palaeontologica Polonica* **56**(3):535–552. doi:10.4202/app.2010.0003
- 396 Gilmore, Charles W. 1936. Osteology of *Apatosaurus* with special reference to specimens in the  
397 Carnegie Museum. *Memoirs of the Carnegie Museum* **11**:175–300 and plates XXI–XXXIV.
- 398 Hatcher, Jonathan B. 1901. *Diplodocus* (Marsh): its osteology, taxonomy and probable habits,  
399 with a restoration of the skeleton. *Memoirs of the Carnegie Museum* **1**:1–63 and plates I–XIII.
- 400 Hatcher, J. B. 1903. Osteology of *Haplocanthosaurus* with description of a new species, and  
401 remarks on the probable habits of the Sauropoda and the age and origin of the *Atlantosaurus*  
402 beds; additional remarks on *Diplodocus*. *Memoirs of the Carnegie Museum* **2**:1–75 and plates  
403 I–VI.

- 404 Huene, Friedrich von. 1932. Die fossile Reptile-Ordnung Saurischia, ihre Entwicklung und  
405 Geschichte. *Monographien zur Geologie und Palaeontologie* (Serie 1) **4**:1–361.
- 406 Ibiricu, Lucio M., Matthew C. Lamanna, Rubén D.F. Martínez, Gabriel A. Casal, Ignacio A.  
407 Cerda, Gastón Martínez and Leonardo Salgado. 2017. A novel form of postcranial skeletal  
408 pneumaticity in a sauropod dinosaur: Implications for the paleobiology of Rebbachisauridae.  
409 *Acta Palaeontologica Polonica* **62(2)**:221–236. doi:10.4202/app.00316.2016
- 410 Lavocat, René J. M. 1954. Sur les Dinosauriens du continental intercalaire des Kem-Kem de la  
411 Daoura. [On the dinosaurs of the Continental Intercalaire of the Kem Kem of the Daoura].  
412 *Comptes Rendus 19th International Geological Congress* **1952(1)**:65–68. English translation  
413 by Matthew C. Lamanna provided by the *Polyglot Paleontologist* at  
414 [http://paleoglot.org/files/Lavocat\\_54.pdf](http://paleoglot.org/files/Lavocat_54.pdf)
- 415 Lydekker, Richard. 1893. On a sauropodous dinosaurian vertebra from the Wealden of Hastings.  
416 *Quarterly Journal of the Geological Society, London* **49**:276–280.
- 417 Mallison, Heinrich, and Oliver Wings. 2014. Photogrammetry in paleontology — a practical  
418 guide. *Journal of Paleontological Techniques* **12**:1–31.
- 419 Mannion, Philip D. 2009. A rebbachisaurid sauropod from the Lower Cretaceous of the Isle of  
420 Wight, England. *Cretaceous Research* **30**:521–526. doi:10.1016/j.cretres.2008.09.005
- 421 Mannion, Philip D. 2010. A revision of the sauropod dinosaur genus '*Bothriospondylus*' with a  
422 redescription of the type material of the middle jurassic form '*B. madagascariensis*'.  
423 *Palaeontology* **53(2)**:277–296. doi:10.1111/j.1475-4983.2009.00919.x
- 424 Mannion, Philip D., Paul Upchurch and Stephen Hutt. 2011. New rebbachisaurid (Dinosauria:  
425 Sauropoda) material from the Wessex Formation (Barremian, Early Cretaceous), Isle of Wight,  
426 United Kingdom. *Cretaceous Research* **32(6)**:774–780. doi:10.1016/j.cretres.2011.05.005
- 427 Marsh, Othniel C. 1878. Principal characters of American Jurassic dinosaurs, part I. *American*  
428 *Journal of Science* (Series 3) **16**:411–416.
- 429 Naish, Darren, and David M. Martill. 2001. Saurischian dinosaurs I: Sauropods. pp. 185–241 in:  
430 Martill, David M., and Darren Naish (eds.). *Dinosaurs of the Isle of Wight*. Palaeontological  
431 Association, London.
- 432 Osborn, Henry Fairfield, and Charles C. Mook. 1921. *Camarasaurus*, *Amphicoelias* and other  
433 sauropods of Cope. *Memoirs of the American Museum of Natural History*, new series  
434 **3(3)**:247–387, and plates LX–LXXXV.
- 435 Owen, Richard. 1842. Report on British fossil reptiles, Part II. *Reports of the British Association*  
436 *for the Advancement of Science* **11**:60–204.
- 437 Riggs, Elmer S. 1903. *Brachiosaurus altithorax*, the largest known dinosaur. *American Journal*  
438 *of Science* **15(4)**:299–306.
- 439 Riggs, Elmer S. 1904. Structure and relationships of opisthocoelian dinosaurs. Part II, the  
440 Brachiosauridae. *Field Columbian Museum, Geological Series* **2(6)**:229–247, plus plates  
441 LXXI–LXXXV.
- 442 Russell, Dale A. 1996. Isolated dinosaur bones from the Middle Cretaceous of the Tafilalet,  
443 Morocco. *Bulletin du Muséum National d'Histoire Naturelle*, 4ème série – section C: Sciences  
444 de la Terre, Paléontologie, Géologie, Minéralogie **18(2–3)**:349–402.

- 445 Seeley, Harry G. 1888. On the classification of the fossil animals commonly named Dinosauria.  
446 *Proceedings of the Royal Society of London* **43**:165–171.
- 447 Sereno, Paul C., and Jeffrey A. Wilson. 2005. Structure and evolution of a sauropod tooth battery.  
448 pp. 157–177 in: Wilson, Jeffrey A., and Kristina Curry-Rogers (eds.), *The Sauropods:*  
449 *Evolution and Paleobiology*. University of California Press, Berkeley.
- 450 Sereno, Paul C, Jeffrey A. Wilson, Lawrence A. Witmer, John A. Whitlock, Abdoulaye Maga,  
451 Oumarou Ide, Timothy A. Rowe. 2007. *Nigersaurus taqueti* (On-line), Digital Morphology.  
452 Accessed November 14, 2017 at  
453 [http://digimorph.org/specimens/Nigersaurus\\_taqueti/dorsal\\_vertebra/](http://digimorph.org/specimens/Nigersaurus_taqueti/dorsal_vertebra/)
- 454 Sereno, Paul C., Allison L. Beck, Didier. B. Dutheil, Hans C. E. Larsson, Gabrielle. H. Lyon,  
455 Bourahima Moussa, Rudyard W. Sadleir, Christian A. Sidor, David J. Varricchio, Gregory P.  
456 Wilson and Jeffrey A. Wilson. 1999. Cretaceous sauropods from the Sahara and the uneven  
457 rate of skeletal evolution among dinosaurs. *Science* **282**:1342–1347.
- 458 Taylor, Michael P. 2015. Is *Xenoposeidon* a rebbachisaur? *Sauropod Vertebra Picture of the Week*,  
459 July 14, 2015. Accessed November 15, 2017 at [https://svpow.com/2015/07/14/is-](https://svpow.com/2015/07/14/is-xenoposeidon-a-rebbachisaur/)  
460 [xenoposeidon-a-rebbachisaur/](https://svpow.com/2015/07/14/is-xenoposeidon-a-rebbachisaur/)
- 461 Taylor, Michael P., and Darren Naish. 2007. An unusual new neosauropod dinosaur from the  
462 Lower Cretaceous Hastings Beds Group of East Sussex, England. *Palaeontology* **50(6)**:1547–  
463 1564. doi: 10.1111/j.1475-4983.2007.00728.x
- 464 Wedel, Mathew J., Richard L. Cifelli and R. Kent Sanders. 2000. Osteology, paleobiology, and  
465 relationships of the sauropod dinosaur *Sauroposeidon*. *Acta Palaeontologica Polonica*  
466 **45(4)**:343–388.
- 467 Wilson, Jeffrey A. 2012. New vertebral laminae and patterns of serial variation in vertebral  
468 laminae of sauropod dinosaurs. *Contributions from the Museum of Paleontology, University of*  
469 *Michigan* **32(7)**:91–110. <http://hdl.handle.net/2027.42/92460>
- 470 Wilson, Jeffrey A., and Ronan Allain. 2015. Osteology of *Rebbachisaurus garasbae* Lavocat,  
471 1954, a diplodocoid (Dinosauria, Sauropoda) from the early Late Cretaceous-aged Kem Kem  
472 beds of southeastern Morocco. *Journal of Vertebrate Paleontology* **35(4)**:e1000701.  
473 doi:10.1080/02724634.2014.1000701

## 474 **Figure Captions**

- 475 **Figure 1.** NHMUK R2095, the holotype and only vertebra of *Xenoposeidon proneneukos*, shown  
476 from all six cardinal directions. Top row: **A.** dorsal view, with anterior to the left. Middle row, left  
477 to right: **B.** anterior, **C.** left lateral, **D.** posterior and **E.** right lateral view. Bottom row: **F.** ventral  
478 view, with anterior to the left. Scale bar = 200 mm.
- 479 **Figure 2.** NHMUK R2095, the holotype and only vertebra of *Xenoposeidon proneneukos*, in left  
480 lateral view, with interpretative drawing. **A.** The incorrect interpretation of the laminae from  
481 Taylor and Naish (2017:figure 4A), with identifying captions greyed out since they are largely  
482 incorrect. **B.** The revised interpretation of the same laminae, based on the similar arrangement in  
483 *Rebbachisaurus garasbae*. Scale bar = 200 mm.
- 484 **Figure 3.** Centra and neural arches of posterior dorsal vertebrae from two rebbachisaurid  
485 sauropods (not to scale), highlighting the distinctive “M” shape formed by laminae high on the

486 neural arch. **A.** NHMUK R2095, the holotype and only vertebra of *Xenoposeidon proneneukos*.  
487 **B.** MNHN MRS 1958, a posterior dorsal vertebra from the holotype specimen of *Rebbachisaurus*  
488 *garasbae*.

489 **Figure 4.** NHMUK R2095, the holotype and only vertebra of *Xenoposeidon proneneukos*, in left  
490 lateral view, interpreted as a rebbachisaurid. This interpretation is modelled primarily on MNHN  
491 MRS 1958, a posterior dorsal vertebra from the holotype specimen of *Rebbachisaurus garasbae*.  
492 The CPOL passes through a sheetlike PCDL, as in *Rebbachisaurus*; but the lateral CPRL forms a  
493 cross-shaped junction with the ACPL, each of these laminae equally interrupting the trajectory of  
494 the other. Abbreviations as used in the text. Scale bar = 200 mm.

495 **Figure 5.** NHMUK R2095, the holotype and only vertebra of *Xenoposeidon proneneukos*, in left  
496 anteroventrolateral view, highlighting the three sets of laminae related to the prezygapophyses.  
497 The trajectories of the medial CPRLs (which emerge from the neural arch pedicels) and the  
498 lateral CPRLs (which intersect with the APCLs) indicate the approximate position of the  
499 prezygapophyses. The additional arched laminae form the margins of the large, teardrop-shaped  
500 CPRF, but meet at a position some way below and posterior to the presumed location of the  
501 prezygapophyseal facets. Breakage of both medial CPRLs and the left ACPL and PCDL is  
502 indicated by cross-hatching. Note that, from this perspective, the lateral CPRL appears to turn a  
503 corner where it intersects with the ACPL, such that the posteroventral portion of the lateral CPRL  
504 appears contiguous with the dorsal portion of the ACPL. This is an illusion brought about by the  
505 eminence at the point of intersection. As always, this is much easier to see in three dimensions.  
506 Abbreviations as used in the text.


## 507 **Supplementary Files**

508 **Supplementary file 1.** Three-dimensional surface model (11 million polygons) of NHMUK  
509 R2095, the holotype and only vertebra of *Xenoposeidon proneneukos*. A 3D polygon mesh file  
510 was created by Heinrich Mallison in Agisoft Photoscan Pro version 1.3.0 (agisoft.com), from 95  
511 high resolution digital photographs by the author. All 95 images aligned, and resulted in a dense  
512 point cloud at maximum resolution of 20,900,043 points and 44,871,128 polygons. Scaling was  
513 based on a single 10 cm scale bar created from a high quality scale bar placed in the pictures with  
514 the specimen. Available from <https://doi.org/10.6084/m9.figshare.5605612.v2>

# Figure 1

NHMUK R2095, the holotype and only vertebra of *Xenoposeidon proneneukos*, shown from all six cardinal directions.

**Figure 1.** NHMUK R2095, the holotype and only vertebra of *Xenoposeidon proneneukos*, shown from all six cardinal directions. Top row: **A.** dorsal view, with anterior to the left. Middle row, left to right: **B.** anterior, **C.** left lateral, **D.** posterior and **E.** right lateral view. Bottom row: **F.** ventral view, with anterior to the left. Scale bar = 200 mm.






## Figure 2

Figure 2. NHMUK R2095, the holotype and only vertebra of *Xenoposeidon proneneukos*, in left lateral view, with interpretative drawing.


**Figure 2.** NHMUK R2095, the holotype and only vertebra of *Xenoposeidon proneneukos*, in left lateral view, with interpretative drawing. **A.** The incorrect interpretation of the laminae from Taylor and Naish (2017:figure 4A), with identifying captions greyed out since they are largely incorrect. **B.** The revised interpretation of the same laminae, based on the similar arrangement in *Rebbachisaurus garasbae*. Scale bar = 200 mm.



## Figure 3

Figure 3. Centra and neural arches of posterior dorsal vertebrae from two rebbachisaurid sauropods (not to scale), highlighting the distinctive “M” shape formed by laminae high on the neural arch.


**Figure 3.** Centra and neural arches of posterior dorsal vertebrae from two rebbachisaurid sauropods (not to scale), highlighting the distinctive “M” shape formed by laminae high on the neural arch. **A.** NHMUK R2095, the holotype and only vertebra of *Xenoposeidon proneneukos*. **B.** MNHN MRS 1958, a posterior dorsal vertebra from the holotype specimen of *Rebbachisaurus garasbae*.



## Figure 4

Figure 4. NHMUK R2095, the holotype and only vertebra of *Xenoposeidon proneneukos*, in left lateral view, interpreted as a rebbachisaurid.

**Figure 4.** NHMUK R2095, the holotype and only vertebra of *Xenoposeidon proneneukos*, in left lateral view, interpreted as a rebbachisaurid. This interpretation is modelled primarily on MNHN MRS 1958, a posterior dorsal vertebra from the holotype specimen of *Rebbachisaurus garasbae*. The CPOL passes through a sheetlike PCDL, as in *Rebbachisaurus*; but the lateral CPRL forms a cross-shaped junction with the ACPL, each of these laminae equally interrupting the trajectory of the other. Abbreviations as used in the text. Scale bar = 200 mm.



## Figure 5

Figure 5. NHMUK R2095, the holotype and only vertebra of *Xenoposeidon proneneukos*, in left anteroventrolateral view, highlighting the three sets of laminae related to the prezygapophyses.

**Figure 5.** NHMUK R2095, the holotype and only vertebra of *Xenoposeidon proneneukos*, in left anteroventrolateral view, highlighting the three sets of laminae related to the prezygapophyses. The trajectories of the medial CPRLs (which emerge from the neural arch pedicels) and the lateral CPRLs (which intersect with the APCLs) indicate the approximate position of the prezygapophyses. The additional arched laminae form the margins of the large, teardrop-shaped CPRF, but meet at a position some way below and posterior to the presumed location of the prezygapophyseal facets. Breakage of both medial CPRLs and the left ACPL and PCDL is indicated by cross-hatching. Note that, from this perspective, the lateral CPRL appears to turn a corner where it intersects with the ACPL, such that the posteroventral portion of the lateral CPRL appears contiguous with the dorsal portion of the ACPL. This is an illusion brought about by the eminence at the point of intersection. As always, this is much easier to see in three dimensions. Abbreviations as used in the text.

