

# RAMANUJAN, ROBIN, THE RIEMANN HYPOTHESIS, AND RECENT RESULTS

JONATHAN SONDOW

`jsondow@alumni.princeton.edu`

RAMANUJAN 125 CONFERENCE November 2012

arXiv:1211.6944v2 [math.HO] 29 Mar 2013



Srinivasa RAMANUJAN (1887–1920)

**Abstract.** The first part of the paper is on the mathematics of my topic, which is the work of Ramanujan and Robin on the Riemann Hypothesis (RH). The second part is on the history.

## 1. MATHEMATICS

**Definition.** The *sum-of-divisors function*  $\sigma$  is defined by

$$\sigma(n) := \sum_{d|n} d = n \sum_{d|n} \frac{1}{d}.$$

For example,  $\sigma(6) = 12$ .

In 1913, Grönwall found the maximal order of  $\sigma$ .

**Grönwall's Theorem.** *The function*

$$G(n) := \frac{\sigma(n)}{n \log \log n} \quad (n > 1)$$

*satisfies*

$$\limsup_{n \rightarrow \infty} G(n) = e^\gamma = 1.78107 \dots,$$

*where  $\gamma = 0.57721 \dots$  is Euler's constant.*



Thomas Hakon GRÖNWALL (1877–1932)

Grönwall's proof uses:

**Mertens's Theorem.** *If  $p$  denotes a prime, then*

$$\lim_{x \rightarrow \infty} \frac{1}{\log x} \prod_{p \leq x} \frac{p}{p-1} = e^{\gamma}.$$



Franz MERTENS (1840–1927)

**Ramanujan's Theorem.** *If RH is true, then for  $n_0$  large enough,*

$$n > n_0 \implies G(n) < e^\gamma.$$

Here is an excerpt from his proof.

Ramanujan: ... *assume that ...  $s > 0$  ... if  $p$  is the largest prime not greater than  $x$ , then*


$$\begin{aligned} & \frac{\log 2}{2^s - 1} + \frac{\log 3}{3^s - 1} + \frac{\log 5}{5^s - 1} + \cdots + \frac{\log p}{p^s - 1} - C \\ &= \int_2^{\vartheta(x)} \frac{dx}{x^s - 1} - s \int_2^x \frac{x - \vartheta(x)}{x^{1-s}(x^s - 1)^2} dx + O(x^{-s}(\log x)^4). \end{aligned}$$

*But it is known that*

$$x - \vartheta(x) = \sqrt{x} + x^{\frac{1}{3}} + \sum \frac{x^\rho}{\rho} - \sum \frac{x^{\frac{1}{2}\rho}}{\rho} + O\{x^{\frac{1}{5}}\}$$

*where  $\rho$  is a complex root of  $\zeta(s)$ . ...*

The last equation is a variant of the classical explicit formula in prime number theory. This shows “explicitly” (pun intended) how Ramanujan used RH in his proof.



Georg Friedrich Bernhard RIEMANN (1826–1866)

**Robin's Theorem.**  *$RH$  is true if and only if*  
$$n > 5040 (= 7!) \implies G(n) < e^\gamma.$$



Guy ROBIN

# GRANDES VALEURS DE LA FONCTION SOMME DES DIVISEURS ET HYPOTHÈSE DE RIEMANN

G. ROBIN

---

RÉSUMÉ. — Soit  $\sigma$  la fonction  $\sigma(n) = \sum_{d|n} d$ . Le but de cet article est de prouver que si l'hypothèse de Riemann est vraie, on a  $\sigma(n) < e^\gamma n \log \log n$  pour  $n \geq 5041$ . ( $\gamma$  est la constante d'Euler). Si l'hypothèse de Riemann est fausse, on montre aussi qu'il existe une infinité de  $n$  tels que  $\sigma(n) > e^\gamma n \log \log n$  et  $\forall n \geq 3$ ,  $\sigma(n) < e^\gamma n \log \log n + 0,6483 n / \log \log n$ .

ABSTRACT. — Let be the function  $\sigma(n) = \sum_{d|n} d$ . The aim of this paper is to prove that under Riemann's hypothesis,  $\sigma(n) < e^\gamma n \log \log n$  for  $n \geq 5041$ . ( $\gamma$  is Euler's constant). If Riemann's hypothesis is false it is shown that for infinitely many  $n$ ,  $\sigma(n) > e^\gamma n \log \log n$  and  $\forall n \geq 3$ ,  $\sigma(n) < e^\gamma n \log \log n + 0,6483 n / \log \log n$ .

Robin's paper on  $\sigma$  and RH,  
Journal de Mathématiques Pures et Appliquées, 1984



The sum-of-divisors function  $\sigma$  and Euler's totient function  $\phi$ , defined as

$$\phi(n) := \sum_{\substack{1 \leq k \leq n \\ (k,n)=1}} 1 = n \prod_{p|n} \left(1 - \frac{1}{p}\right),$$

are related by the inequalities

$$\frac{6}{\pi^2} < \frac{\sigma(n)}{n} \cdot \frac{\phi(n)}{n} < 1,$$

which hold for  $n > 1$ . They follow from the identity

$$\frac{\sigma(n)}{n} \cdot \frac{\phi(n)}{n} = \prod_{1 < p^e \parallel n} \left(1 - \frac{1}{p^{e+1}}\right).$$

Robin used ideas from a result on the  $\phi$  function proved by his thesis advisor Nicolas in 1983.

**Nicolas's Theorem.**  *$RH$  is true if and only if*

$$p \text{ prime} \implies \frac{\phi(p\#)}{p\# / \log \log p\#} < e^{-\gamma},$$

where  $p\# := 2 \times 3 \times 5 \times \cdots \times p$  denotes a primorial.



Jean-Louis NICOLAS

Nicolas in turn used Landau's 1903 result on the minimal order of the totient function.

**Landau's Theorem.** *The  $\phi$  function satisfies*

$$\liminf_{n \rightarrow \infty} \frac{\phi(n)}{n / \log \log n} = e^{-\gamma} = 0.56145 \dots$$



Edmund LANDAU (1877–1938)

## RECENT RESULTS

**Definition.** An integer  $N > 1$  is a *GA1 number* if  $N$  is composite and  $G(N) \geq G(N/p)$  for all prime factors  $p$ . Call  $N$  a *GA2 number* if  $G(N) \geq G(aN)$  for all multiples  $aN$ .

### **Caveney, Nicolas, and S.'s Theorems.**

1. *RH is true if and only if 4 is the only number that is both GA1 and GA2.*
2. *A GA2 number  $N > 5040$  exists if and only if RH is false, in which case  $N$  is even and  $> 10^{8576}$ .*



Geoffrey CAVENEY

## 2. HISTORY

My story begins in 1915 when Ramanujan published the first part of his dissertation “Highly Composite Numbers” (HCN).

**Papers Published in the Proceedings of the London Mathematical Society**  
**S. Ramanujan**  
**Highly Composite Numbers**  
Proc. London Math. Soc. (1915) s2\_14(1): 347-409 doi:10.1112/plms/s2\_14.1.347

Part 1 of HCN by Ramanujan,  
Proceedings of the London Mathematical Society,  
1915

Ramanujan: *I define a highly composite number as a number whose number of divisors exceeds that of all its predecessors.*


His thesis was written at Trinity College, University of Cambridge, where his advisors were Hardy and Littlewood.



Godfrey Harold HARDY (1877–1947)



John Edensor LITTLEWOOD (1885–1977)



Ramanujan (center) at his degree ceremony, 1916



In 1944, Erdős published a paper “On highly composite and similar numbers” with Alaoglu.

Erdős (in “Ramanujan and I”): *Ramanujan had a very long manuscript on highly composite numbers but some of it was not published due to a paper shortage during the First World War.*




Paul ERDŐS (1913–1996)

Erdős: *Hardy rather liked this paper but perhaps not unjustly called it nice but in the backwaters of mathematics.*

Hardy: *Even Ramanujan could not make highly composite numbers interesting.*

Dyson: *Hardy said this to discourage me from working on H. C. numbers myself. I think he was right.*



Freeman DYSON

In 1982 Rankin published a paper on “Ramanujan’s manuscripts and notebooks”. Rankin quoted Hardy’s 1930 letter to Watson in which Hardy mentioned “the suppressed part of HCN”.

Rankin: *The most substantial manuscript consists of approximately 30 pages of HCN carrying on from where the published paper stops.*

By a curious coincidence, 1981–1982 is also the year when Séminaire Delange-Pisot-Poitou published an exposition of Robin’s Theorem, in which he improved on Ramanujan’s Theorem without ever having heard of it!

Berndt: *It is doubtful that Rankin took notice of Robin’s paper. I definitely did not.*

Thus Rankin and Berndt on the English-American side, and Nicolas and Robin on the French side, were not communicating.



Robert Alexander RANKIN (1915–2001)



Bruce Carl BERNDT holding Ramanujan's slate

Berndt: *After I began to edit Ramanujan's notebooks, I wrote Trinity College in 1978 for a copy of the notes that Watson and Wilson made in their efforts to edit the notebooks. I also decided to write for copies of all the Ramanujan material that was in the Trinity College Library. Included in their shipment to me was the completion of Ramanujan's paper on highly composite numbers. I put all of this on display during the Ramanujan centenary meeting at Illinois in June, 1987.*

Nicolas: *I keep a very strong souvenir of the conference organised in Urbana-Champaign in 1987 for the one hundred anniversary of Ramanujan. It is there that I discovered the hidden part of “Highly Composite Numbers”.*

*What I have not written is that there was an error of calculus in Ramanujan’s manuscript which prevented him from seeing Robin’s Theorem. Soon after discovering the hidden part, I read it and saw the difference between Ramanujan’s result and Robin’s one. Of course, I would have bet that the error was in Robin’s paper, but after recalculating it several times and asking Robin to check, it turned out that there was an error of sign in what Ramanujan had written.*



Ramanujan's Theorem was not explicitly stated by him in HCN. Nicolas and Robin formulated it for him in Note 71 of their annotated and corrected version of HCN Part 2.

Nicolas and Robin: *It follows from [the corrected version of Ramanujan's formula] (382) that under the Riemann hypothesis, and for  $n_0$  large enough,*

$$n > n_0 \implies \sigma(n)/n < e^\gamma \log \log n.$$

*It has been shown in [22] that the above relation with  $n_0 = 5040$  is equivalent to the Riemann hypothesis.*

Here [22] is Robin's paper, which he published three years *before* hearing of Ramanujan's Theorem in 1987. However, a reader of Note 71 who neglects to look up [22] in the References is left with the misimpression that the proof "that the above relation with  $n_0 = 5040$  is equivalent to the Riemann hypothesis" was given *after* "the above relation" became known!



In 1993, HCN Part 2 was submitted to Proceedings of the London Mathematical Society, which had published Part 1 in 1915. The paper was accepted, but could not be published, because Trinity College did not own the rights to Ramanujan's papers and was not able to obtain permission from his widow, Janaki.



S. Janaki Ammal (1899–1994), Mrs. Ramanujan

Janaki passed away in 1994, and the paper was eventually published by Alladi in the first volume of his newly-founded Ramanujan Journal.

THE RAMANUJAN JOURNAL, Vol. 1, No. 2 (1997), 119-153.

## Highly composite numbers

by  
SRINIVASA RAMANUJAN

Annotated by  
**JEAN-LOUIS NICOLAS**  
*Institut Girard Desargues, UPRES-A-5028, Mathématiques, Bâtiment 101, Université Claude Bernard (Lyon 1), F-69622 Villeurbanne cédex, France*  
[jlnicola@frcpn11.in2p3.fr](mailto:jlnicola@frcpn11.in2p3.fr)

and  
**GUY ROBIN**  
*UPRES-A-6090, Théorie des nombres, calcul formel et optimisation, Université de Limoges, F-87060 Limoges cédex, France*  
[robin@cict.fr](mailto:robin@cict.fr)

*Received December 15, 1995; Accepted April 11, 1996*

**Abstract.** In 1915, the London Mathematical Society published in its Proceedings a paper of S. Ramanujan entitled "Highly Composite Numbers." But it was not the whole work on the subject, and in "The Lost Notebook and Other Unpublished Papers," one can find a manuscript, handwritten by Ramanujan, which is the continuation of the paper published by the London Mathematical Society.

This paper is the typed version of the above mentioned manuscript with some notes, mainly explaining the link between the work of Ramanujan and works published after 1915 on the subject.

Part 2 of HCN by Ramanujan,  
annotated by Nicolas and Robin,  
The Ramanujan Journal, 1997



Krishnaswami ALLADI,  
founder of The Ramanujan Journal

Here my story ends. If it has offended anyone, I apologize.



S. with Ramjee RAGHAVAN, Ramanujan's grand-nephew, by chance (!) my seatmate on the first leg of my flight to the Ramanujan 125 Conference in FL