

The Blue Lurker WOCS 14020: A Long-Period Post-Common-Envelope Binary in M67 Originating from a Merger in a Triple System

2 EMILY M. LEINER,^{1,2} NATALIE M. GOSNELL,³ AARON M. GELLER,⁴ MENG SUN,² ROBERT D. MATHIEU,⁵ AND ALISON SILLS⁶

3
4 ¹*Department of Physics, Illinois Institute of Technology, 3101 South Dearborn St., Chicago, IL 60616, USA*

5 ²*Center for Interdisciplinary Exploration and Research in Astrophysics (CIERA), Northwestern University, 1800 Sherman Ave.,*
6 *Evanston, IL 60201, USA*

7 ³*Department of Physics, Colorado College, 14 E. Cache La Poudre St., Colorado Springs, CO 80903, USA*

8 ⁴*Center for Interdisciplinary Exploration and Research in Astrophysics (CIERA) and Department of Physics and Astronomy,*
9 *Northwestern University, 1800 Sherman Ave., Evanston, IL 60201, USA*

10 ⁵*Department of Astronomy, University of Wisconsin - Madison, 475 N. Charter Street, Madison WI 53726*

11 ⁶*Department of Physics and Astronomy, McMaster University, 1280 Main Street West, Hamilton, ON, L8S 3X6, CANADA*

12 ABSTRACT

13 We present *Hubble Space Telescope* far-ultraviolet (FUV) spectra of a blue-lurker–white-dwarf (BL-
14 WD) binary system in the 4 Gyr open cluster M67. We fit the FUV spectrum of the WD, determining
15 it is a C/O WD with a mass of $0.72_{-0.04}^{+0.05} M_{\odot}$ and a cooling age of ~ 400 Myr. This requires a
16 WD progenitor of $\sim 3 M_{\odot}$, significantly larger than the current cluster turnoff mass of $1.3 M_{\odot}$. We
17 suggest the WD progenitor star formed recently via the merger of two stars near the turnoff of the
18 cluster. In this scenario, the original progenitor system was a hierarchical triple consisting of a close,
19 near-equal-mass inner binary, with a tertiary companion with an orbit of a few thousand days. The
20 WD is descended from the merged inner binary, and the original tertiary is now the observed BL.
21 The likely formation scenario involves a common envelope while the WD progenitor is on the AGB,
22 and thus the observed orbital period of 359 days requires an efficient common envelope ejection. The
23 rapid rotation of the BL indicates it accreted some material during its evolution, perhaps via a wind
24 prior to the common envelope. This system will likely undergo a second common envelope in the
25 future, and thus could result in a short-period double WD binary or merger of a $0.72 M_{\odot}$ C/O WD
26 and a $0.38 M_{\odot}$ Helium WD, making this a potential progenitor of an interesting transient such as a
27 sub-Chandrasekhar Type Ia supernova.

28 1. INTRODUCTION

29 Blue lurkers (BLs) are stars that appear in color-magnitude diagrams to be normal main sequence stars (see Figure 1),
30 but they have shorter rotation periods than expected. For instance, the solar-like stars in the old (4 Gyr) open cluster
31 M67 rotate with $P_{rot} \sim 20 - 30$ days. Leiner et al. (2019) detect 11 BLs in M67 with $P_{rot} \leq 8$ days. To explain this
32 rapid rotation, Leiner et al. (2019) hypothesize BLs have been spun up via stellar mergers, collisions, or mass transfer
33 in binary systems. BLs are therefore thought to be the lower-luminosity counterparts to the more well-known blue
34 straggler stars, which are found in open and globular clusters brighter than the main-sequence turnoff (Figure 1). Like
35 the blue stragglers, BLs may be formed via mass transfer from a giant companion (McCrea 1964; Chen & Han 2008),
36 stellar mergers (Perets & Fabrycky 2009), or stellar collisions during dynamical encounters (Knigge et al. 2009; Leigh
37 & Sills 2011).

38 An observational test of the mass-transfer hypothesis for blue straggler formation is to search for white dwarf (WD)
39 companions to blue straggler stars. Studies of the old (6 Gyr) open cluster NGC 188 used *HST* UV photometry
40 and spectroscopy to identify WD companions to blue stragglers (Gosnell et al. 2014, 2015, 2019), finding that $\sim 1/3$
41 of the blue stragglers had detectable hot WD companions. Further, these studies suggested that 2/3 of the blue
42 straggler population likely had WD companions, as some WDs would be too cool and old to be detected. Numerous

UV detections of WD companions to blue stragglers have now been claimed in other clusters using far-UV imaging (e.g. Saketh et al. 2024; Jadhav et al. 2019; Sindhu et al. 2019, Panthi et al. 2022, 2024 and references therein). Direct detection of UV flux from WD companions demonstrates definitively that many blue stragglers form via transfer from a giant companion.

Nine et al. (2023) applied this same technique to the BL population of M67 identified in Leiner et al. (2019), using *HST* UV photometry to look for UV excesses indicative of WD companions. They detected hot, young (< 900 Myr) WDs in two of the BL binary systems (WOCS 3001 and WOCS 14020), confirming that at least $\sim 20\%$ of the BLs have been spun up to rapid rotation rates via mass transfer from a giant binary. The true fraction of mass transfer formation is likely larger as older, cooler WD companions are too faint to be photometrically detected. Jadhav et al. (2019) also report UV excesses to M67 BLs WOCS 3001 and possibly WOCS 9005 using *AstroSat/UVIT*, though the WOCS 9005 detection was not confirmed by Nine et al. (2023). The detection of WD companions to some BLs solidifies the hypothesis that BLs are lower-luminosity, lower mass analogs to the blue straggler stars that blend photometrically with typical main sequence stars, and that some formed via mass transfer from a red giant branch (RGB) or asymptotic giant branch (AGB) donor stars.

While many blue stragglers, BLs and related systems are now known to have WD companions, and thus to have formed via mass transfer of some kind, very few have the detailed stellar and orbital parameters needed to infer detailed formation histories. WD masses and ages, in particular, are essential constraints because they define the evolutionary state of the donor star at the end of mass transfer, and the time since mass transfer occurred. These parameters can only be ascertained from eclipsing or self lensing binaries (e.g. Kawahara et al. 2018) or from fitting FUV WD spectra (e.g. Landsman et al. 1997) and therefore only a few blue stragglers have well characterized formation histories (Landsman et al. 1997; Brogaard et al. 2018; Gosnell et al. 2019; Sun et al. 2021; Sun & Mathieu 2023), and no blue lurkers have yet had a detailed formation history proposed.

Post-mass-transfer blue stragglers and BLs have orbital periods of $10^2 - 10^3$ days. These long orbital periods challenge the often used assumption in population synthesis models that mass transfer from more massive giant stars on to less massive main-sequence accretors should be unstable (Hjellming & Webbink 1987), leading to a common envelope and orbital inspiral to form short-period binaries with orbital periods of just a few days. The wide orbital periods of the BLs may support recent theoretical models that suggest stable mass transfer may occur onto lower-mass accretors than canonically predicted (see, for example Temmink et al. 2023; Ge et al. 2020; Pavlovskii & Ivanova 2015; Passy et al. 2012; Woods & Ivanova 2011). Alternatively, it may be that BLs are not the result of standard stable mass transfer, but instead form via another pathway such as via wind accretion. Even minimal accretion via a wind ($\Delta M < 0.1 M_{\odot}$) has been shown to potentially spin accreting stars up to velocities approaching break up (Sun et al. 2024), potentially explaining the observed rapid rotation in BLs.

In the few cases where the formation histories of blue stragglers have been investigated in detail, the formation pathways include both quite conservative and highly non-conservative mass transfer on the RGB (Landsman et al. 1997; Sun et al. 2021), and a combination of Roche lobe overflow on the RGB/AGB and wind mass transfer (Sun & Mathieu 2023). Thus, formation paths of blue stragglers and blue lurkers appear to be varied, and more detailed case studies are needed to better understand the range of formation scenarios and formation physics.

Here we seek to understand the formation pathway of a BL-WD binary in M67, WOCS 14020, by constraining the WD mass and cooling age from its far-UV spectrum in order to re-construct the evolutionary history of the system. In Section 2 we describe the targeted BL-WD binary, WOCS 14020. In Section 3 we present the observation and spectral analysis technique we use to determine a WD mass and cooling age for this target. These masses and ages inform the possible mass transfer history of this system, which we explore with some modeling in Section 4. We discuss the implications of our findings (Section 5) and conclude with a summary of our results (Section 6).

2. TARGET AND OBSERVATIONS

2.1. WOCS 14020

WOCS 14020 is a binary star system in the 4 Gyr, solar-metallicity cluster M67 (Geller et al. 2015; Leiner et al. 2019; Geller et al. 2021). It is a single-lined spectroscopic binary dominated by the light from the BL primary star. Given this system's position in the CMD (Figure 1), the BL star is consistent with a $\sim 1.0 M_{\odot}$ main-sequence star. The BL primary was found in Leiner et al. (2019) to have a rotation period of 4.4 days, unusually fast for a solar-like star at an age of 4 Gyr, which have expected rotation periods > 20 days. Gyrochronology models (Angus et al. 2019) predict an age of ~ 300 Myr given this rotation period. Nine et al. (2023) analyzed *HST* FUV photometry of WOCS

		Comment
M	$\sim 1.05 M_{\odot}$	Estimated from evolutionary track fit to photometry
T_{eff}	5990^{+60}_{-100} K	SED fit in Nine et al. (2023)
M_G	4.76	Gaia DR3 absolute magnitude using $d = 816$ pc and $E(B - V) = 0.041$
$(bp - rp)_0$	0.78	Gaia DR3 color with $E(B-V) = 0.041$ and Wang & Chen (2019) extinction law
P_{rot}	4.4 days	From Leiner et al. (2019)
P_{orb}	358.9 days	From Leiner et al. (2019)
ecc	0.23	From Leiner et al. (2019)
$f(m)$	2.38×10^{-3}	From Geller et al. (2021)
$M_{2,\text{min}}$	$0.15 M_{\odot}$	Calculated from $f(m)$ (Leiner et al. 2019)
$M_{2,\text{orbit}}$	$0.35 M_{\odot}$	Predicted from P_{orb} & $ecc.$ using Rappaport et al. (1995)
Age (Myr)	~ 300	Using Angus et al. (2019) gyrochronology models

Table 1. Stellar and Orbital Properties of WOCS 14020

14020, detecting a FUV excess consistent with a hot WD companion. The photometry is consistent with a temperature for the WD of $\sim 11,000 - 13,000$ K, implying a time since mass-transfer formation of $\sim 290 - 540$ Myr, consistent with the gyrochronology age. Assuming a primary mass of $1.05 M_{\odot}$, the binary mass function also yields a very low minimum secondary mass ($0.15 M_{\odot}$), consistent with a WD secondary. We summarize the system properties of the BL primary and the orbital parameters of the binary system in Table 1.

2.2. HST Spectroscopy

WOCS 14020 was observed by *HST* Cosmic Origins Spectrograph (COS) for GO program 17134 over five separate visits of two orbits each. It was observed in TIME-TAG mode through the Primary Science Aperture (PSA) using the G140L grating with a central wavelength of 1105Å. This region covers the Lyman- α wings and provides the wide wavelength coverage necessary to fit WD atmosphere models in this region. To increase the signal-to-noise of the final spectrum, we co-added the MAST-reduced spectrum for each visit and binned the resulting spectrum to a new wavelength resolution of 3.0Å. The spectrum is dereddened using `pysynphot` tools assuming $E(B - V) = 0.041$ ([Taylor 2007](#)). Geocoronal emission lines are evident in the spectrum and are masked by hand from the analysis.

3. SPECTRAL FITTING

To constrain the WD parameters, we fit the reduced and combined COS spectrum with WD atmosphere models ([Koester 2010](#)) using the MCMC tool `emcee` ([Foreman-Mackey et al. 2013](#)). An MCMC approach best captures the inherent degeneracy between $\log g$ and T_{eff} for WD atmosphere fits. The normalization of the fitted WD atmosphere scales with r^2/d^2 , where r is the WD radius and d is the distance to the system. We adopt the M67 cluster distance from [Stello et al. \(2016\)](#) of 816 ± 11 pc. The radius of a WD depends on both the surface gravity and temperature, as well as the core composition of the WD. Rather than make an assumption about the core composition, we include a third “core-picking” parameter that chooses whether to fit using a He-core composition (corresponding to $\log g < 7.7$, [Althaus et al. 2013](#)) or a CO-core composition (corresponding to $\log g \geq 7.7$, [Holberg & Bergeron 2006](#); [Tremblay et al. 2011](#)). We apply flat priors on $\log g$ and T_{eff} ranging from 6.0–9.0 and 10000–18000 K, respectively, and a flat prior across the core-picking parameter.

We ran 300 walkers for 30000 steps with a thinning factor of 10 and a burn-in of 1000, for 6×10^5 final samples. The resulting autocorrelation times for $\log g$, T_{eff} , and the core-picking parameter are 8.6, 8.2, and 6.3, respectively, demonstrating that the model fits are stable. The model fitting posteriors are shown in Figure 2, with best fit values (using 16th and 84th percentiles) of $\log g = 8.17^{+0.09}_{-0.06}$ cm s^{-2} and $T_{\text{eff}} = 13400^{+240}_{-160}$ K. From the fitted $\log g$ and T_{eff} values, we calculate the corresponding core mass and cooling age ranges by interpolating standard WD mass-radius relationships ([Holberg & Bergeron 2006](#); [Tremblay et al. 2011](#); [Althaus et al. 2013](#)). The resulting derived core mass and cooling age are $M_{\text{WD}} = 0.72^{+0.05}_{-0.04} M_{\odot}$ and 390^{+40}_{-30} Myr.

A He-core WD is completely eliminated as a reasonable fit to the spectrum. The `emcee` results definitively point not only to a C/O-core WD, but a *surprisingly massive* C/O-core WD of approximately $0.7 M_{\odot}$ given its very young age. We note there is a small portion of the posterior distribution (approximately 2%) that allows for a moderate WD





Figure 1. Color-magnitude diagram of M67 highlighting *Gaia* DR3 proper-motion members (gray), blue stragglers (blue), BLs (orange), and our target, the BL WOCS 14020 (red square). Memberships are determined as described in Leiner & Geller (2021)


128 mass of $0.55 M_{\odot}$, as can be seen on the left edge of the posterior distributions in Figure 2, but the vast majority of
 129 the posterior is consistent with a massive WD companion.

130 4. FORMATION PATHWAY

131 The measured mass of the WD companion star, $0.72^{+0.05}_{-0.04} M_{\odot}$ is significantly larger than the predicted WD mass
 132 that would result from end-state evolution of any turnoff mass star at the age of M67 ($M_{\text{WD}} \sim 0.55 M_{\odot}$). The WD
 133 age we determine is ~ 400 Myr, and the age of M67 is generally found to be between 3.5-4.2 Gyr (Stello et al. 2016;
 134 Sarajedini et al. 2009; Barnes et al. 2016). Based on this age range, the mass transfer would have occurred when a
 135 typical giant star in M67 would have been $\lesssim 1.6 M_{\odot}$. According to the COSMIC (Breivik et al. 2020, a rapid population
 136 synthesis code based on pre-calculated BSE stellar evolutionary tracks), this would yield a final WD mass of < 0.57
 137 M_{\odot} . Therefore, our measured WD mass is significantly more massive than a WD that could have been produced



(a)



(b)

Figure 2. (a) Binned COS spectrum for WOCS 14020, overlaid with 100 random draws from the posterior distribution from the WD atmosphere fits. The light blue data points correspond to Earth-shine emission and are not included in the fitting routine. (b) Posterior probability distributions for the WD atmosphere fits to the COS spectrum, with the median, 16–84% (approximating 1 sigma), and 2.5–97.5% (approximating 2 sigma) percentile values marked as indicated in the legend. The $\log g$ and T_{eff} values are fit directly and the corresponding WD mass and age are calculated using CO-core mass-radius relationships and cooling times.

by a typical single star in M67. In fact, given standard single-star stellar evolution, such a massive WD requires a progenitor with a mass close to $3 M_{\odot}$. Such a massive star would have a main sequence lifetime of < 1 Gyr, and therefore $0.72 M_{\odot}$ WDs in M67 should have cooling ages of more than 3 Gyr. The observed cooling age of ~ 400 Myr for the WD in WOCs 14020 necessitates that a $3 M_{\odot}$ star was created in the cluster quite recently, likely through a binary merger or interaction. We note that Williams et al. (2018) (see also Canton et al. 2021) detect a single WD with a similar mass in M67, and also infer that it descended from a stellar merger. The detection of this similarly overmassive white dwarf supports our interpretation that this is the result of a merger.


In Figure 3, we show COSMIC model grids of potential progenitor binary systems. We use default COSMIC parameters on a grid of solar metallicity main sequence binaries with initially circular orbits between 100 and 10000 days. We vary the initial primary mass from 2.6 to $3.9 M_{\odot}$, and use a fixed secondary mass of $1.0 M_{\odot}$. The colors indicate the final remnant WD mass, with colors chosen to emphasize the regions where the WD mass falls within the 16th–84th percentile region of our WD mass determination, what we will refer to as our confidence interval.

According to panel (a), the progenitor of this WD must be larger than $2.8 M_{\odot}$ to produce a WD with a mass within our confidence interval. Progenitors with masses approaching $4.0 M_{\odot}$ are also possible if they are in binaries with $P_{\text{orb}} \lesssim 10^3$ days, where the core growth will be truncated by the onset of mass transfer. However, given that the turnoff mass at the time of formation was less than $1.6 M_{\odot}$, the maximum mass of a star produced via a merger of two stars in the cluster would be $3.2 M_{\odot}$. Therefore, we suggest the likely progenitor of this WD was a $2.8 - 3.2 M_{\odot}$ star, approximately twice the turnoff mass. Within this mass range, the progenitor binary would have had an initial orbital period of a few thousand days or more to produce a WD of the observed mass.


In panels (b) and (c) of Figure 3, we show a similar plot to (a), but plotting the final orbital periods of the binary grid after mass transfer. We show the observed orbital period of the BL-WD binary with a dashed black line. These models indicate that mass-transfer in this system would have been unstable, resulting in a common envelope (CE) that shrinks the orbital period. Initial orbital periods of a few thousand days will shrink to periods of several hundred days or less, depending on the efficiency of the CE ejection, matching the observed 359 day period of WOCs 14020. For these grids we use COSMIC default parameters, except we vary the common envelope (CE) efficiency α . This α parameter describes the fraction of the orbital energy that can be used to unbind the giant envelope (see Hurley et al. 2002 and Breivik et al. 2020 for a complete description of the CE treatment in BSE/COSMIC). The CE λ parameter, which relates to the stellar envelope binding energy, is set using the default COSMIC option of adopting the value from Claeys et al. (2014). This CE prescription can produce a binary with the observed orbital period of WOCs 14020 following an episode of CE evolution, provided the CE ejection efficiency is high ($\alpha > 0.8$.) We note, however, that our models assume all binaries start in circular orbits, and stay circularized as they evolve. WOCs 14020 is currently in a moderately eccentric orbit. This eccentricity is difficult to reproduce in models because it is not theoretically understood, which we discuss further in Section 5.2.

Based on this modeling, the observed stellar and orbital properties of the BL-WD binary, and the additional constraints imposed by the star’s membership in M67, we propose this system likely formed via the merger of an inner binary in a hierarchical triple, followed by mass transfer from the merger remnant onto the wide tertiary companion resulting in a CE. This triple may have been primordial, or may have formed dynamically. In more detail:


1. This system begins as a hierarchical triple. The inner binary consists of two stars with near equal masses of $M \sim 1.5 M_{\odot}$ in a short-period binary system. The system has a main-sequence tertiary companion with $M \sim 1.0 M_{\odot}$ in a wide orbit of several thousand days.
2. The inner binary undergoes a merger, perhaps induced by stellar evolution as one of the components begins to evolve off the main sequence, magnetic braking, or due to Kozai-Lidov cycles (Kozai 1962; Lidov 1962) induced by the tertiary companion (e.g. Perets & Fabrycky 2009; Naoz & Fabrycky 2014). The merger remnant is $\sim 3.0 M_{\odot}$ and retains the initial tertiary as a binary companion with an orbital period of $P_{\text{orb}} \sim 3500$ days.
3. About 400 Myr ago, the $\sim 3.0 M_{\odot}$ merger remnant evolves into an AGB star. Near the tip of the AGB, wind mass transfer begins and the original tertiary companion accretes a small amount of material via wind. This spins up the rotation rate of the accretor, yielding the rapid rotation we see in the BL today.
4. Soon a CE is triggered and the binary inspirals. A high common envelope efficiency results in the common envelope being quickly lost from the system. Its initial period of ~ 3500 day shrinks to the currently observed period of 359 days.



(a)



(b) $\alpha = 1.0$



(c) $\alpha = 0.8$

Figure 3. (top) COSMIC model grid showing the initial primary mass and initial orbital period of our post-merger binary. Colors indicates the mass of the C/O WD that emerges after the final CE and emphasize final WD masses that fall within our mass confidence interval (0.68-0.78 M_{\odot}). (bottom) The progenitor masses and orbital periods of the binary system *after* evolution through mass-transfer or a CE. On the left we show models using $\alpha = 1.0$ and on the right we show models using $\alpha = 0.8$. The 359-day orbital period of WOCS 14020 is indicated with the vertical dashed line.

- 188 5. The WD begins to cool and the BL spins down via magnetic braking.
- 189 6. About 400 Myr later, we now observe the BL-WD binary as it is today: a 1.05 M_{\odot} BL primary with a 0.72 M_{\odot}
- 190 WD secondary in a binary with $P_{\text{orb}} = 359$ days.

191 5. DISCUSSION

192 We have argued above that WOCS 14020 is a remarkable example of the complexity of evolution in a triple system.

193 We propose that its evolution has included a merger of an inner binary in a hierarchical triple, followed by a common

194 envelope between the merger product and the original tertiary. Here we discuss several implications of this case-study

195 in triple star evolution.

196 5.1. Common Envelope Evolution

The current orbital period of WOCS 14020 indicates that the system must have evolved through a CE, as the separation is much too small to host a giant star with a $0.72 M_{\odot}$ core. Given a progenitor donor star of $2.8 - 3.2 M_{\odot}$ and the current WD mass of $0.72 M_{\odot}$, the orbital period at the onset of the CE must have been a few thousand days (Figure 3 a). The CE ejection efficiency must have been quite high to result in the observed amount of orbital decay. COSMIC models require a value of the CE α of 0.8–1.0 to reproduce the observed post-CE orbital period of this system (Figure 3 (b) and (c)). This CE efficiency is a large uncertainty in binary evolution, with some studies arguing that more inefficient envelope ejections with $\alpha \sim 0.2 - 0.3$ better match observed the characteristics of the post-CE MS-WD population (see, for example Zorotovic et al. 2010; Toonen & Nelemans 2013; Camacho et al. 2014 or double WD systems Scherbak & Fuller 2023), while others require $\alpha \sim 1.0$ or even larger (De Marco et al. 2011; Sun & Arras 2018). Most previous investigations have looked at post-CE binaries with very short orbital periods of a few days or WDs with low-mass M-dwarf companions to constrain α (e.g. Zorotovic et al. 2010; Scherbak & Fuller 2023), generally finding a much lower efficiency of $\sim 0.2 \lesssim \alpha \lesssim 0.4$ is required. Our observations offer a rare constraint on outcomes of CE that involve solar-type companions and result in orbital periods of hundreds of days. Some similar systems have recently been discovered that also favor efficient ejection of CEs initiated during the donor’s AGB phase (Belloni et al. 2024; Yamaguchi et al. 2024a), as does the recent discovery of a large population of MS-WD binaries in the field with orbital periods in the range of $10^2 - 10^3$ days (Yamaguchi et al. 2024b). Our observation and analysis also favor this efficient envelope ejection.

5.2. Orbital Eccentricity

Notably, WOCS 14020 has a non-zero eccentricity of $e = 0.23$. Outcomes of stable mass transfer and CE evolution have often been assumed to be circular. However, it is common for blue stragglers, BLs, and related post-mass-transfer binaries such as binary post-AGB, s-processed enhanced stars, and main sequence-WD binaries to have non-zero eccentricities (e.g. Mathieu & Geller 2009; Leiner et al. 2019; Oomen et al. 2018; Escorza et al. 2019; Shahaf et al. 2024; Yamaguchi et al. 2024b). The reasons for this are still uncertain. Some work has proposed that eccentricity pumping during stable mass transfer can occur (Sepinsky et al. 2007, 2009, 2010; Rocha et al. 2024, in prep), which may explain some of these eccentric post-mass-transfer binaries, but WOCS 14020 does not appear to have evolved through stable mass transfer. CE events are generally assumed to result in circularized binary systems, but a growing body of work also questions this assumption. Accretion from a circumbinary disk is a potential mechanism to excite eccentricities in binaries (Dermine et al. 2013), including post-CE binaries with circumbinary disks (Kashi & Soker 2011; Siwek et al. 2023; Valli et al. 2024; Wei et al. 2024). Alternatively, the binary may not actually circularize prior to the CE as is often assumed, and some eccentricity may be preserved after the CE (e.g. Bonačić Marinović et al. 2008; Prust & Chang 2019).

5.3. Mass Accretion and Spin Up

The rapid rotation of the BL argues for some accretion during the system’s evolution. It is often assumed that no accretion occurs on to the secondary star that evolves through CE evolution, but the rapid rotation requires some accretion happened before, during, or after the CE event. Sun et al. (2024) recently showed that even a small amount of wind accretion ($< 0.1 M_{\odot}$) can spin up accretors to critical rotation rates. In our COSMIC model, the proto-BL accretes $\sim 0.1 M_{\odot}$ via a wind prior to the onset of CE, and this may be enough to explain the observed rotation. Nine et al. (2024) recently also surveyed the blue stragglers and BLs in M67 for Barium enhancement, an often used indicator of mass transfer from an AGB companion. They did not find WOCS 14020 to be significantly barium enhanced relative to the main sequence population. This may also support the interpretation that only minimal accretion took place, which was insufficient to significantly increase the Barium surface abundance but was sufficient to spin up the accretor considerably. While we note that only minimal accretion is needed to produce the rapid rotation and observed color-magnitude diagram position of the BL, our observations do not rule out larger accretion amounts, and other accretion mechanisms besides a wind prior to CE onset may also be possible. For example, there could be accretion from a circumbinary disk (Lai & Muñoz 2023) that forms after the CE (Kashi & Soker 2011), which could transfer mass to the BL, spin it up, and lead to the observed orbital eccentricity as noted in the previous section.

5.4. The Importance of Triples

Given what we know about stellar multiplicity and binary statistics, evolutionary paths similar to WOCS 14020 may be fairly common. Around 10% of solar-type stars are found in triple systems (Raghavan et al. 2010), and a

configuration with a short-period near-equal mass inner binary and a wider tertiary is common among these systems. Moe & Di Stefano (2017) find an excess of “twin” binaries among short period ($P < 100$ day) solar-like binaries such that $\sim 30\%$ of short-period binaries have near equal mass ratios ($0.95 < \frac{M_2}{M_1} < 1.0$). A large fraction of close binaries have also been found to have wide tertiary companions; Tokovinin et al. (2006) find the overall triple fraction among close spectroscopic binaries ($P_{\text{orb}} < 30$ days) is $63 \pm 5\%$, and that rises to 96% among the closest binaries with $P_{\text{orb}} < 3$ days. Therefore, mergers of inner binaries in triples should often form stars that are nearly twice the turnoff mass of a cluster, and these merger products will often have companions they will interact with later in their evolution.

Growing observational evidence also points to the general importance of triples in stellar evolution. Heintz et al. (2022) analyze the *Gaia* sample of wide double WD binaries, which reveals a large fraction ($\sim 20\%$) of systems include a WD that resulted from a merger, and thus originated in a triple system. Shariat et al. (2024) also argue that in $\sim 40\%$ of wide double WD binaries, the more massive WD is a merger product, and thus these systems are descended from triples. Further, they find that 20-25% of blue stragglers may form from evolution in triples.

In M67, Leiner et al. (2016) detected an overmassive giant with a binary companion, S1237. This system consists of a red giant primary that is an asteroseismic outlier for the cluster. Asteroseismic analysis yields a mass of $2.9 \pm 0.2 M_{\odot}$, more than twice the turnoff mass of M67, and indicates the star is likely a core helium burning giant. This overmassive giant has a binary companion in a 697.8 day orbit, which seems to be located near the cluster turnoff or perhaps in the blue straggler region. The most likely formation scenario for S1237 is quite similar to WOCS 14020: the overmassive giant likely resulted from the merger of a close, near-equal-mass inner binary in a hierarchical triple. The current binary companion would then have previously been a wide tertiary. The giant in S1237 is expected to form a WD in the near future (~ 100 Myr), at which point the system should be observed as a blue straggler- or BL-WD binary. Another blue straggler system in M67, S1082, is currently a triple system containing two blue stragglers, and must have formed from multiple mergers or collisions given the system’s large combined mass (Sandquist et al. 2003; Leigh & Sills 2011). Descendants of triple systems are thus not rare amongst the known post-mass-transfer population of M67; at least one appears to be produced every few hundred Myr.

Theoretical work has also highlighted the potential importance of creating blue stragglers, double WD binaries, barium stars, and other post-interaction systems via triples (e.g. Perets & Fabrycky 2009; Toonen et al. 2022; Gao et al. 2023). This reflects a growing theoretical consensus that considering triple systems is vital to understanding the full breadth of stellar evolution pathways and outcomes. WOCS 14020 is among the best observationally-characterized examples of a post-interaction triple system to date, and thus is a much needed test case for future models of evolution in triple systems.

5.5. Future Evolution of WOCS 14020

Intriguingly, WOCS 14020 could be a potential Type Ia supernova progenitor via a sub-Chandrasekhar double detonation (Liu et al. 2023b), or the progenitor of a calcium-rich transient (Kasliwal et al. 2012; Jacobson-Galán et al. 2021; Morán-Fraile et al. 2024). Based on past detailed case studies of open cluster blue stragglers forming through mass transfer (Sun et al. 2021; Sun & Mathieu 2023), in approximately 5 Gyr, the BL will evolve into a giant star and begin to interact with the C/O WD companion. It is unlikely WOCS 14020 will be disrupted by a dynamical encounter within the next 5 Gyr (Leigh & Sills 2011), so it is probable that the system will evolve towards a double WD system of some kind. Most likely, a CE will occur, leaving behind a close double WD binary consisting of the $0.72 M_{\odot}$ C/O WD and a $0.38 M_{\odot}$ He WD that is the remnant of the current BL. Depending on the CE efficiency, this double WD binary may be close enough to interact, with the larger He WD accreting onto the C/O WD. The outcomes of such sub-Chandrasekhar mass mergers and interactions are still not well understood, but recent models suggest a variety of interesting transients may result.

In a detailed binary evolution study, Wong & Bildsten (2023) simulated a close He WD transferring mass onto a C/O WD and discussed the necessary conditions for triggering a Type Ia supernova event. One of the requirements is to have a high-entropy He WD, which could be formed from a CE event. In general, our system configuration aligns well with Wong & Bildsten (2023) for achieving a successful explosion event. Furthermore, it has recently been suggested that C/O WD with $M < 1.0 M_{\odot}$ can undergo a detonation by accreting stably or unstably from a He WD companion (Shen et al. 2024), resulting in a normal Type Ia supernova. Some Type Ia supernova have been observed with similar C/O WD progenitor masses (see De et al. 2019; Liu et al. 2023a). Mergers between C/O and He WD have also been suggested as potential sources of calcium-rich supernovae (e.g. Morán-Fraile et al. 2024). The evolutionary history of

WOCS 14020 underscores that triple evolution may be a source of late-time transients, including double detonation Type Ia supernova and calcium-rich transients.

6. SUMMARY AND CONCLUSION

WOCS 14020 is a BL-WD binary in the open cluster M67. It has a 359-day orbital period with a moderate eccentricity of $e = 0.23$. We fit the Lyman- α region of a *HST*/COS far-UV spectrum of the WD in this system. From this fit we determine $\log g = 8.17^{+0.09}_{-0.06}$ cm s $^{-2}$ and $T_{\text{eff}} = 13400^{+240}_{-160}$ K for this WD. This corresponds to a C/O WD with $M_{\text{WD}} = 0.72^{+0.05}_{-0.04} M_{\odot}$ and a cooling age of 390^{+40}_{-30} Myr. The mass is significantly larger than expected for a typical WD of this age in this cluster, and we argue that the progenitor star was a $2.8 - 3.2 M_{\odot}$ merger product. We suggest a formation scenario in which this BL-WD binary is descended from a hierarchical triple system in the cluster, which undergoes first a merger of the inner binary, followed by mass transfer onto the outer tertiary that ends with a CE that shrinks the orbit from ~ 3500 days to its currently observed 359 day period. We note that:

- The rapid rotation of the BL indicates it accreted at least a small amount of material during the interaction, possibly via a wind prior to CE and/or accretion from a post-CE disk.
- The non-circular orbit indicates eccentricity is either somehow maintained through the CE evolution or excited after the CE.
- the current orbital period of 359 days indicates an efficient envelope ejection ($\alpha \gtrsim 0.8$).

In the future, this system will likely undergo a second CE when the BL evolves into a red giant and overflows its Roche lobe. Ultimately, this may form a close double WD binary or yield a He WD-C/O WD merger. The story of WOCS 14020 thus underscores the complexities of evolution in triple star systems, and highlights that triple evolution is an important channel creating blue stragglers, double WD binaries, explosive transients, and other important astrophysical objects that result from stellar interactions. WOCS 14020 is the first blue lurker system for which an evolutionary path has been illuminated in detail, a result made possible because its stellar and orbital parameters have been exceptionally well-constrained by observations. This case study offers rare and intriguing insights about the outcome of interactions in a triple star system.

ACKNOWLEDGMENTS

This paper is based on observations made with the NASA/ESA *Hubble Space Telescope*, obtained at the Space Telescope Science Institute, which is operated by the Association of Universities for Research in Astronomy, Inc., under NASA contract NAS 5-26555. These observations are associated with program #17134 and supported by NASA HST-GO-17134. This work has made use of data from the European Space Agency (ESA) mission *Gaia* (<https://www.cosmos.esa.int/gaia>), processed by the *Gaia* Data Processing and Analysis Consortium (DPAC, <https://www.cosmos.esa.int/web/gaia/dpac/consortium>). Funding for the DPAC has been provided by national institutions, in particular the institutions participating in the *Gaia* Multilateral Agreement. E.L. is additionally supported by funding from Illinois Institute of Technology. M.S. acknowledges the support from the GBMF8477 grant (PI: Vicky Kalogera) and thanks Chang Liu and Sunny Wong for discussions on double detonation type Ia supernovae. R.M. acknowledges funding from the Wisconsin Alumni Research Foundation. A.S. is supported by the Natural Sciences and Engineering Research Council of Canada.

Software: COSMIC (Breivik et al. 2020), emcee (Foreman-Mackey et al. 2013) numpy (Harris et al. 2020), Matplotlib (Hunter 2007), astropy (Astropy Collaboration et al. 2013)

All the *HST* data used in this paper can be found in MAST: <http://dx.doi.org/10.17909/twcy-vq55>

REFERENCES

- | | |
|--|--|
| <p>336 Althaus, L. G., Miller Bertolami, M. M., & Córscico, A. H.
337 2013, A&A, 557, A19</p> <p>338 Angus, R., Morton, T. D., Foreman-Mackey, D., et al.
339 2019, AJ, 158, 173</p> | <p>340 Astropy Collaboration, Robitaille, T. P., Tollerud, E. J.,
341 et al. 2013, A&A, 558, A33</p> <p>342 Barnes, S. A., Weingrill, J., Fritzewski, D., Strassmeier,
343 K. G., & Platais, I. 2016, ApJ, 823, 16</p> |
|--|--|

- 344 Belloni, D., Schreiber, M. R., & Zorotovic, M. 2024, arXiv
345 e-prints, arXiv:2401.17510
- 346 Bonačić Marinović, A. A., Glebbeek, E., & Pols, O. R.
347 2008, *A&A*, 480, 797
- 348 Breivik, K., Coughlin, S., Zevin, M., et al. 2020, *ApJ*, 898,
349 71
- 350 Brogaard, K., Christiansen, S. M., Grundahl, F., et al.
351 2018, *MNRAS*, 481, 5062
- 352 Camacho, J., Torres, S., García-Berro, E., et al. 2014,
353 *A&A*, 566, A86
- 354 Canton, P. A., Williams, K. A., Kilic, M., & Bolte, M.
355 2021, *AJ*, 161, 169
- 356 Chen, X., & Han, Z. 2008, *MNRAS*, 387, 1416
- 357 Claeys, J. S. W., Pols, O. R., Izzard, R. G., Vink, J., &
358 Verbunt, F. W. M. 2014, *A&A*, 563, A83
- 359 De, K., Kasliwal, M. M., Polin, A., et al. 2019, *ApJL*, 873,
360 L18
- 361 De Marco, O., Passy, J.-C., Moe, M., et al. 2011, *MNRAS*,
362 411, 2277
- 363 Dermine, T., Izzard, R. G., Jorissen, A., & Van Winckel, H.
364 2013, *A&A*, 551, A50
- 365 Escorza, A., Karinkuzhi, D., Jorissen, A., et al. 2019, *A&A*,
366 626, A128
- 367 Foreman-Mackey, D., Hogg, D. W., Lang, D., & Goodman,
368 J. 2013, *PASP*, 125, 306
- 369 Gao, Y., Toonen, S., & Leigh, N. 2023, *MNRAS*, 518, 526
- 370 Ge, H., Webbink, R. F., Chen, X., & Han, Z. 2020, *ApJ*,
371 899, 132
- 372 Geller, A. M., Latham, D. W., & Mathieu, R. D. 2015, *AJ*,
373 150, 97
- 374 Geller, A. M., Mathieu, R. D., Latham, D. W., et al. 2021,
375 *AJ*, 161, 190
- 376 Gosnell, N. M., Leiner, E. M., Mathieu, R. D., et al. 2019,
377 *ApJ*, 885, 45
- 378 Gosnell, N. M., Mathieu, R. D., Geller, A. M., et al. 2015,
379 *ApJ*, 814, 163
- 380 —. 2014, *ApJL*, 783, L8
- 381 Harris, C. R., Millman, K. J., van der Walt, S. J., et al.
382 2020, *Nature*, 585, 357–362
- 383 Heintz, T. M., Hermes, J. J., El-Badry, K., et al. 2022,
384 *ApJ*, 934, 148
- 385 Hjellming, M. S., & Webbink, R. F. 1987, *ApJ*, 318, 794
- 386 Holberg, J. B., & Bergeron, P. 2006, *AJ*, 132, 1221
- 387 Hunter, J. D. 2007, *Computing in Science & Engineering*, 9,
388 90
- 389 Hurley, J. R., Tout, C. A., & Pols, O. R. 2002, *MNRAS*,
390 329, 897
- 391 Jacobson-Galán, W. V., Margutti, R., Kilpatrick, C. D.,
392 et al. 2021, *ApJL*, 908, L32
- 393 Jadhav, V. V., Sindhu, N., & Subramaniam, A. 2019, *ApJ*,
394 886, 13
- 395 Kashi, A., & Soker, N. 2011, *MNRAS*, 417, 1466
- 396 Kasliwal, M. M., Kulkarni, S. R., Gal-Yam, A., et al. 2012,
397 *ApJ*, 755, 161
- 398 Kawahara, H., Masuda, K., MacLeod, M., et al. 2018,
399 *ArXiv e-prints*, arXiv:1801.07874
- 400 Knigge, C., Leigh, N., & Sills, A. 2009, *Nature*, 457, 288
- 401 Koester, D. 2010, *Mem. Soc. Astron. Italiana*, 81, 921
- 402 Kozai, Y. 1962, *AJ*, 67, 591
- 403 Lai, D., & Muñoz, D. J. 2023, *ARA&A*, 61, 517
- 404 Landsman, W., Aparicio, J., Bergeron, P., Di Stefano, R.,
405 & Stecher, T. P. 1997, *ApJL*, 481, L93
- 406 Leigh, N., & Sills, A. 2011, *MNRAS*, 410, 2370
- 407 Leiner, E., Mathieu, R. D., Stello, D., Vanderburg, A., &
408 Sandquist, E. 2016, *ApJL*, 832, L13
- 409 Leiner, E., Mathieu, R. D., Vanderburg, A., Gosnell, N. M.,
410 & Smith, J. C. 2019, *ApJ*, 881, 47
- 411 Leiner, E. M., & Geller, A. 2021, *ApJ*, 908, 229
- 412 Lidov, M. L. 1962, *Planet. Space Sci.*, 9, 719
- 413 Liu, C., Miller, A. A., Polin, A., et al. 2023a, *ApJ*, 946, 83
- 414 Liu, Z.-W., Röpkke, F. K., & Han, Z. 2023b, *Research in*
415 *Astronomy and Astrophysics*, 23, 082001
- 416 Mathieu, R. D., & Geller, A. M. 2009, *Nature*, 462, 1032
- 417 McCrea, W. H. 1964, *MNRAS*, 128, 147
- 418 Moe, M., & Di Stefano, R. 2017, *ApJS*, 230, 15
- 419 Morán-Fraile, J., Holas, A., Röpkke, F. K., Pakmor, R., &
420 Schneider, F. R. N. 2024, *A&A*, 683, A44
- 421 Naoz, S., & Fabrycky, D. C. 2014, *ApJ*, 793, 137
- 422 Nine, A. C., Mathieu, R. D., Gosnell, N. M., & Leiner,
423 E. M. 2023, *ApJ*, 944, 145
- 424 Nine, A. C., Mathieu, R. D., Schuler, S. C., & Milliman,
425 K. E. 2024, *ApJ*, 970, 187
- 426 Oomen, G.-M., Van Winckel, H., Pols, O., et al. 2018,
427 *A&A*, 620, A85
- 428 Panthi, A., Vaidya, K., Jadhav, V., et al. 2022, *MNRAS*,
429 516, 5318
- 430 Panthi, A., Vaidya, K., Vernekar, N., et al. 2024, *MNRAS*,
431 527, 8325
- 432 Passy, J.-C., Herwig, F., & Paxton, B. 2012, *ApJ*, 760, 90
- 433 Pavlovskii, K., & Ivanova, N. 2015, *MNRAS*, 449, 4415
- 434 Perets, H. B., & Fabrycky, D. C. 2009, *ApJ*, 697, 1048
- 435 Prust, L. J., & Chang, P. 2019, *MNRAS*, 486, 5809
- 436 Raghavan, D., McAlister, H. A., Henry, T. J., et al. 2010,
437 *ApJS*, 190, 1
- 438 Rappaport, S., Podsiadlowski, P., Joss, P. C., Di Stefano,
439 R., & Han, Z. 1995, *MNRAS*, 273, 731
- 440 Saketh, P., Panthi, A., & Vaidya, K. 2024, *AJ*, 168, 97
- 441 Sandquist, E. L., Latham, D. W., Shetrone, M. D., &
442 Milone, A. A. E. 2003, *AJ*, 125, 810

- 443 Sarajedini, A., Dotter, A., & Kirkpatrick, A. 2009, *ApJ*,
444 698, 1872
- 445 Scherbak, P., & Fuller, J. 2023, *MNRAS*, 518, 3966
- 446 Sepinsky, J. F., Willems, B., Kalogera, V., & Rasio, F. A.
447 2007, *ApJ*, 667, 1170
- 448 —. 2009, *ApJ*, 702, 1387
- 449 —. 2010, *ApJ*, 724, 546
- 450 Shahaf, S., Hallakoun, N., Mazeh, T., et al. 2024, *MNRAS*,
451 529, 3729
- 452 Shariat, C., Naoz, S., El-Badry, K., et al. 2024, *arXiv*
453 e-prints, [arXiv:2407.06257](https://arxiv.org/abs/2407.06257)
- 454 Shen, K. J., Boos, S. J., & Townsley, D. M. 2024, *arXiv*
455 e-prints, [arXiv:2405.19417](https://arxiv.org/abs/2405.19417)
- 456 Sindhu, N., Subramaniam, A., Jadhav, V. V., et al. 2019,
457 *ApJ*, 882, 43
- 458 Siwek, M., Weinberger, R., & Hernquist, L. 2023, *MNRAS*,
459 522, 2707
- 460 Stello, D., Vanderburg, A., Casagrande, L., et al. 2016,
461 *ApJ*, 832, 133
- 462 Sun, M., & Arras, P. 2018, *ApJ*, 858, 14
- 463 Sun, M., Levina, S., Gossage, S., et al. 2024, *ApJ*, 969, 8
- 464 Sun, M., & Mathieu, R. D. 2023, *ApJ*, 944, 89
- 465 Sun, M., Mathieu, R. D., Leiner, E. M., & Townsend,
466 R. H. D. 2021, *ApJ*, 908, 7
- 467 Taylor, B. J. 2007, *AJ*, 133, 370
- 468 Temmink, K. D., Pols, O. R., Justham, S., Istrate, A. G., &
469 Toonen, S. 2023, *A&A*, 669, A45
- 470 Tokovinin, A., Thomas, S., Sterzik, M., & Udry, S. 2006,
471 *A&A*, 450, 681
- 472 Toonen, S., Boekholt, T. C. N., & Portegies Zwart, S. 2022,
473 *A&A*, 661, A61
- 474 Toonen, S., & Nelemans, G. 2013, *A&A*, 557, A87
- 475 Tremblay, P.-E., Bergeron, P., & Gianninas, A. 2011, *ApJ*,
476 730, 128
- 477 Valli, R., Tiede, C., Vigna-Gómez, A., et al. 2024, *A&A*,
478 688, A128
- 479 Wang, S., & Chen, X. 2019, *ApJ*, 877, 116
- 480 Wei, D., Schneider, F. R. N., Podsiadlowski, P., et al. 2024,
481 *A&A*, 688, A87
- 482 Williams, K. A., Canton, P. A., Bellini, A., et al. 2018,
483 *ApJ*, 867, 62
- 484 Wong, T. L. S., & Bildsten, L. 2023, *ApJ*, 951, 28
- 485 Woods, T. E., & Ivanova, N. 2011, *ApJL*, 739, L48
- 486 Yamaguchi, N., El-Badry, K., Ciardi, D. R., et al. 2024a,
487 *PASP*, 136, 074201
- 488 Yamaguchi, N., El-Badry, K., Rees, N. R., et al. 2024b,
489 *PASP*, 136, 084202
- 490 Zorotovic, M., Schreiber, M. R., Gänsicke, B. T., & Nebot
491 Gómez-Morán, A. 2010, *A&A*, 520, A86